B. Initial clinical assessment Refer women who are Women with pain and/or bleeding, or haemodynamically unstable, or in atypical symptoms suggestive of whom there is significant concern early pregnancy problems [see about the degree of pain or recommendation 11] bleeding, directly to A&E. (exit pathway) Healthcare professional conducts clinical history taking, physical examination and urine pregnancy test (unless very recent pregnancy test result reported by the woman) Positive pregnancy test Negative pregnancy test (exit pathway) No pelvic tenderness, cervical Pelvic tenderness. cervical motion tenderness, abdominal motion tenderness, or pain and tenderness or signs of intraabdominal tenderness examination abdominal bleeding Women with a pregnancy of Women with: less than 6 weeks gestation pain or who are bleeding but not in pain a pregnancy of 6 weeks gestation or more or a pregnancy of uncertain gestation Use expectant management. Advise women to repeat a urine pregnancy test after 7-10 days and to return if their symptoms continue or worsen or the test is positive. $(!)(\mathfrak{P})(\lozenge)$ Refer to an early pregnancy assessment service (or out-ofhours gynaecology service if Refer immediately for further early pregnancy the investigation at an early assessment service is not pregnancy assessment service Positive urine Negative available). The urgency of this (or out-of-hours gynaecology pregnancy test, pregnancy test referral depends on the service if the early pregnancy or continuing or (exit pathway) clinical situation. (see section assessment service is worsening C) available). (see section C) symptoms