



## Myofibrillar Myopathy – RETIRED CHAPTER, FOR HISTORICAL REFERENCE ONLY

Duygu Selcen, MD<sup>1</sup> and Andrew G Engel, MD<sup>1</sup>

Created: January 28, 2005; Updated: October 29, 2012.

### Summary

**NOTE: THIS PUBLICATION HAS BEEN RETIRED. THIS ARCHIVAL VERSION IS FOR HISTORICAL REFERENCE ONLY, AND THE INFORMATION MAY BE OUT OF DATE.**

### Clinical characteristics

Myofibrillar myopathy is characterized by slowly progressive weakness that can involve both proximal and distal muscles. Distal muscle weakness is present in about 80% of individuals and is more pronounced than proximal weakness in about 25%. A minority of individuals experience sensory symptoms, muscle stiffness, aching, or cramps. Peripheral neuropathy is present in about 20% of affected individuals. Overt cardiomyopathy is present in 15%-30%.

### Diagnosis/testing

The diagnosis of myofibrillar myopathy is based on clinical findings, electromyography (EMG), nerve conduction studies, and, most importantly, muscle histology. To date, the genetic basis of myofibrillar myopathy has been elucidated in only about 50% of cases. Pathogenic variants have been identified in *DES*, the gene encoding desmin; *CRYAB*, encoding alpha-crystallin B chain; *MYOT*, encoding myotilin; *LDB3* (*ZASP*), encoding LIM domain-binding protein 3; *FLNC*, encoding filamin-C, and *BAG3*, encoding BAG family molecular chaperone regulator 3. Recently, pathogenic variants in *FHL1*, encoding four and a half LIM domains protein 1, and *DNAJB6*, encoding DnaJ homolog subfamily B member 6 were also identified.

### Management

*Treatment of manifestations:* Consider pacemaker and implantable cardioverter defibrillator (ICD) in individuals with arrhythmia and/or cardiac conduction defects; consider cardiac transplantation in individuals with progressive or life-threatening cardiomyopathy; respiratory support (continuous or bilevel positive airway pressure), initially at night and later in the daytime, in individuals with hypercapnea and other signs of incipient

respiratory failure; range-of-motion physical therapy and assistive devices for those with advanced muscle weakness.

*Other:* The role of strengthening exercises has not been defined.

## Genetic counseling

Myofibrillar myopathy is most commonly inherited in an autosomal dominant manner. Exceptions include: X-linked inheritance of *FHL1* pathogenic variants and autosomal recessive inheritance of *CRYAB* pathogenic frameshift variants that lead to premature termination of the translational chain resulting in non-transcription of the mutated protein. The inheritance in some families cannot be determined with confidence. When the pathogenic variant(s) in the family are known, carrier testing and prenatal testing for pregnancies at increased risk is possible.

## GeneReview Scope

Myofibrillar Myopathy: Included Disorders
<ul style="list-style-type: none"> <li>• Alpha-B crystallinopathy</li> <li>• <i>BAG3</i>-related myofibrillar myopathy</li> <li>• Desminopathy</li> <li>• <i>DNAJB6</i>-related myofibrillar myopathy</li> <li>• <i>FHL1</i>-related myofibrillar myopathy</li> <li>• Filaminopathy</li> <li>• Myotilinopathy</li> <li>• Zaspopathy</li> </ul>

For synonyms and outdated names see Nomenclature.

## Diagnosis

### Clinical Diagnosis

The term myofibrillar myopathy refers to a group of genetically distinct disorders linked by common morphologic features observed on muscle histology.

The diagnosis of myofibrillar myopathy rests on the following:


- **History of slowly progressive weakness** accompanied in a smaller proportion of individuals by paresthesias, muscle atrophy, stiffness or aching, cramps, dyspnea, and dysphagia. Physical examination reveals proximal and distal weakness in the majority of individuals. In about one third the weakness is greater proximally than distally; in another third it is greater distally than proximally. Facial weakness is uncommon but can occur. Tendon reflexes are normal or diminished.
- **Electromyography (EMG)** that reveals abnormal electrical irritability (fibrillation potentials, positive sharp waves, complex repetitive discharges, and myotonic discharges) in most individuals. The motor unit potentials show either myopathic features or both myopathic and neurogenic features. Abnormal nerve conduction studies are detected in about 20% of individuals.
- **Muscle histology** that reveals the following combination of findings (see Figure 1):
  - Characteristic alterations in trichromatically stained frozen sections consisting of amorphous, hyaline, or granular material in a variable proportion of the muscle fibers;
  - Sharply circumscribed decreases of oxidative enzyme activity in many abnormal fiber regions;
  - Intense congophilia of many hyaline structures, best observed under rhodamine fluorescence optics; and
  - Small vacuoles in a variable number of fibers

- **Immunocytochemical studies of muscle** that show abnormal ectopic expression of myotilin (90% of abnormal fibers), desmin (75%),  $\alpha$ -B crystallin (75%), dystrophin (70%), and  $\beta$ -amyloid precursor protein (70%)
- **Electron microscopy of muscle** showing progressive myofibrillar degeneration commencing at the Z-disk, disintegration of the sarcomeres, accumulation of degraded filamentous material in pleomorphic hyaline inclusions, and dislocation of membranous organelles and their degradation in autophagic vacuoles [Selcen et al 2004]
- **Other tissues**
  - Peripheral nerve biopsies, in a few descriptions, show accumulation of neurofilaments, neurotubules, and formation of axonal spheroids.
  - Myocardial biopsies show desmin-immunoreactive cytoplasmic inclusions, especially near intercalated disks and interstitial fibrosis.
  - Peripheral nerve pathology [Sabatelli et al 1992] and myocardial pathology [Abraham et al 1998, Arbustini et al 1998, Lohr et al 1998] have been described briefly in a small number of individuals with myofibrillar myopathy.
- **Serum creatine kinase concentration** can be normal or elevated to no greater than seven times the upper limit of normal.

## Molecular Genetic Testing

**Genes.** To date, the genetic basis of myofibrillar myopathy (MFM) has been elucidated in about 50% of kinships. Pathogenic variants have been identified in the following genes:

- *DES*, encoding desmin [Goldfarb et al 1998, Muñoz-Mármol et al 1998, Dalakas et al 2000]
- *CRYAB*, encoding  $\alpha$ -crystallin B chain [Vicart et al 1998, Selcen & Engel 2003]
- *MYOT* (known previously as *TTID*), encoding myotilin [Selcen & Engel 2004]
- *LDB3* (known previously as *ZASP*), encoding LIM domain-binding protein 3 [Selcen & Engel 2005]
- *FLNC*, encoding filamin-C [Vorgerd et al 2005]
- *BAG3*, encoding BAG family molecular chaperone regulator 3 (Bag3) [Selcen et al 2009]
- *FHL1*, encoding four and a half LIM containing protein 1 [Selcen et al 2011]
- *DNAJB6*, encoding DnaJ homolog subfamily B member 6 (heat shock protein 40) [Harms et al 2012; Sarparanta et al 2012; Author, personal observation]



**Figure 1.** Muscle histology observed in myofibrillar myopathy

**Table 1.** Molecular Genetic Testing Used in Myofibrillar Myopathy

Gene <sup>1</sup>	Proportion of MFM Attributed to Mutation of This Gene <sup>2</sup>	Test Method	Variants Detected <sup>3</sup>
<i>DES</i>	7%	Sequence analysis <sup>4</sup>	Sequence variants <sup>5</sup>
		Deletion/duplication analysis <sup>6</sup>	Unknown, none reported <sup>7</sup>
<i>CRYAB</i>	3%	Sequence analysis <sup>4</sup>	Sequence variants
		Deletion/duplication analysis <sup>6</sup>	Unknown; none reported <sup>7</sup>
<i>MYOT</i>	9%	Sequence analysis <sup>4</sup>	Sequence variants
		Deletion/duplication analysis <sup>6</sup>	Unknown, none reported <sup>7</sup>
<i>LDB3</i>	11%	Sequence analysis <sup>4</sup>	Sequence variants
		Sequence analysis of select exons	Sequence variants in exon 6 <sup>8</sup>
		Deletion/duplication analysis <sup>6</sup>	Unknown, none reported <sup>7</sup>
<i>FLNC</i>	3%	Sequence analysis <sup>4</sup>	Sequence variants
		Deletion/duplication analysis <sup>6</sup>	Unknown, none reported <sup>7</sup>
<i>BAG3</i>	5%	Sequence analysis <sup>4</sup>	Sequence variants
		Deletion/duplication analysis <sup>6</sup>	Unknown, none reported <sup>7</sup>
<i>FHL1</i>	3%	Sequence analysis <sup>4</sup>	Sequence variants

Table 1. continued from previous page.

Gene <sup>1</sup>	Proportion of MFM Attributed to Mutation of This Gene <sup>2</sup>	Test Method	Variants Detected <sup>3</sup>
<i>DNAJB6</i>	2%	Sequence analysis <sup>4</sup>	Sequence variants

1. See Table A. Genes and Databases for chromosome locus and protein.

2. Frequencies based on Mayo Clinic MFM cohort

3. See Molecular Genetics for information on allelic variants.

4. Sequence analysis detects variants that are benign, likely benign, of uncertain significance, likely pathogenic, or pathogenic. Pathogenic variants may include small intragenic deletions/insertions and missense, nonsense, and splice site variants; typically, exon or whole-gene deletions/duplications are not detected. For issues to consider in interpretation of sequence analysis results, click [here](#).

5. Expected to identify approximately 99% of pathogenic variants in the coding region.

6. Testing that identifies deletions/duplications not readily detectable by sequence analysis of the coding and flanking intronic regions of genomic DNA; included in the variety of methods that may be used are: quantitative PCR, long-range PCR, multiplex ligation-dependent probe amplification (MLPA), and chromosomal microarray (CMA) that includes this gene/chromosome segment.

7. No large deletions or duplications of *DES*, *MYOT*, *LDB3*, *CRYAB*, *FLNC*, or *BAG3* have been reported to cause myofibrillar myopathy.

8. Exons sequenced may vary among laboratories.

## Testing Strategy

### To confirm/establish the diagnosis in a proband

- Generic diagnosis of an MFM can be made on clinical and histologic grounds.
- The diagnosis of a specific MFM is based on molecular genetic testing:
  - If the proband's symptoms suggest a candidate gene (see Genotype-Phenotype Correlations), perform sequence analysis of that gene first.
  - If no pathogenic variant is identified, perform sequence analysis for the remaining genes (Table 1).
  - At present no clear criteria for testing for deletions/duplications exist.

**Predictive testing** for at-risk asymptomatic adult family members requires prior identification of the pathogenic variant(s) in the family.

**Prenatal diagnosis and preimplantation genetic diagnosis (PGD)** for at-risk pregnancies require prior identification of the pathogenic variant(s) in the family.

## Clinical Characteristics

### Clinical Description

In the Mayo Clinic series of 80 individuals with myofibrillar myopathy (MFM), the age of onset varied from two to 77 years. The age at diagnosis ranged from 11 to 82 years. *BAG3*-related myofibrillar myopathy (Bag3opathy) characteristically presents in the first or second decade of life and is typically fatal. Desminopathies may also present in the first decade of life, usually with cardiomyopathy. However, the majority of MFM presents after age 40 years.

The predominant presenting symptom in MFM is slowly progressive weakness; a minority of individuals experience sensory symptoms, muscle stiffness, aching, or cramps. The weakness can involve both proximal and distal muscles; however, distal muscle weakness is 25% more common than proximal weakness.

Objective clinical signs or EMG findings of peripheral neuropathy are present in approximately 20% of affected individuals, but muscle biopsy studies suggest an even higher frequency of peripheral nerve involvement.

Overt cardiomyopathy can be a presenting manifestation or can appear during the evolution of myofibrillar myopathy in 15%-30% of affected individuals.

A restrictive ventilatory defect can result from respiratory muscle weakness.

Variable expressivity has been observed in kinships with pathogenic variants in *DES*, with some family members showing signs of cardiomyopathy only, some showing signs of both myopathy and cardiomyopathy, and some with reduced penetrance who have signs of neither myopathy nor cardiomyopathy.

## Genotype-Phenotype Correlations

No morphologic features consistently or reliably predict mutation of a given gene.

The only genotype-phenotype correlations detected to date are:

- Cataracts were present in affected members of one of the three reported kinships with pathogenic variants in *CRYAB*, the gene encoding alpha crystallin B chain (also known as  $\alpha$ -B crystallin) [Vicart et al 1998].
- Neuropathy and cardiomyopathy were observed in families with pathogenic variants in *MYOT* [Selcen & Engel 2004] and *BAG3* [Selcen et al 2009].
- Rigid spine has been observed in Bag3opathy [Selcen et al 2009] and *FHL1*-related MFM [Selcen et al 2011].
- Neuropathy and cardiomyopathy also occur in individuals in whom the genetic basis of myofibrillar myopathy has not been established.

## Penetrance

Data are insufficient to draw conclusions about penetrance.

## Anticipation

No convincing evidence of anticipation has been documented.

## Nomenclature

"... The light microscopic features of myofibrillar myopathy were described in the 1970s and 1980s under such names as "myopathy with inclusion bodies" [Nakashima et al 1970], "atypical myopathy with myofibrillar aggregates" [Kinoshita et al 1975], "autosomal dominant cardiomyopathy with inclusions" [Clark et al 1978], "cardioskeletal myopathy with intrasarcoplasmic dense granulofilamentous material" [Fardeau et al 1978], "autosomal dominant spheroid body myopathy" [Goebel et al 1978], "myopathy with sarcoplasmic bodies and desmin filaments" [Edström et al 1980], "congenital myopathy with cytoplasmic bodies" [Wolburg et al 1982], "congenital myopathy with Mallory body-like inclusions" [Fidzianska et al 1983], and "familial cardiomyopathy with subsarcolemmal vermiform deposits" [Calderon et al 1987]. Later, however, it became apparent that some authors described the same pathologic reaction under different names; that more than one pathologic alteration thought to be specific for a single disorder could be present in the same muscle specimen, or even in the same muscle fiber; and that in each instance the pathologic changes involve the Z-disks of the myofibril. Edström et al [1980] noted that some inclusions or material around them reacted for desmin. This generated the names of "desmin storage myopathy" [Horowitz & Schmalbruch 1994] and later "desmin-related myopathies" [Goebel 1997]. In 1996 and 1997, detailed immunocytochemical studies revealed that many proteins, not just desmin, accumulate in the abnormal fibers, and prompted the use of the noncommittal term myofibrillar myopathy [De Bleeker et al 1996, Nakano et al 1997]..."

[Selcen et al 2004; republished with permission of Oxford University Press]



Myofibrillar myopathy has also been referred to as desmin storage myopathy, desmin-related myopathy, or protein surplus myopathy. Because myofibrillar myopathy is genetically heterogeneous and the disease-causing protein or involved gene is known only in a minority of cases, because multiple other proteins besides desmin are also overexpressed in muscle, and because myotilin is not related to desmin, the generic term "myofibrillar myopathy" is the preferred designation until the gene(s) in which pathogenic variants occur are determined. When the disease-associated gene or protein is identified, designations such as desminopathy,  $\alpha$ -B crystallinopathy, myotilinopathy, zaspopathy (Markesbery-Griggs late-onset distal myopathy), filaminopathy, or *BAG3*-related myofibrillar myopathy are appropriate.

## Prevalence

The prevalence of myofibrillar myopathy cannot be estimated at this time.

## Genetically Related (Allelic) Disorders

Cardiomyopathy can occur with each genetically defined type of MFM.

Mutation of *CRYAB* can also cause cataracts without causing myofibrillar myopathy.

Mutation of *MYOT* can also cause a limb girdle-like phenotype that has been designated as LGMD1A [Hauser et al 2000] and also as a distal myopathy [Fischer et al 2006, Pénisson-Besnier et al 2006].

Mutation of *FHL1* is associated with several phenotypes: X-linked childhood-onset reducing body myopathy; X-linked early-onset severe reducing body myopathy; *FHL1*-related X-linked [Emery-Dreifuss muscular dystrophy](#); X-linked myopathy with postural muscle atrophy; X-linked dominant scapuloperoneal myopathy.

Mutation of *DBA/IB6* is associated with limb-girdle muscular dystrophy type 1E.

## Differential Diagnosis

The principle differential diagnoses are late-onset myopathies and especially myopathies with a predominantly distal distribution:

- **Myotonic dystrophy type 1 (DM1)** is a multisystem disorder that affects skeletal and smooth muscle as well as the eye, heart, endocrine system, and central nervous system. The clinical findings, which span a continuum from mild to severe, have been categorized into three somewhat overlapping phenotypes: mild, classic, and congenital. Mild DM1 is characterized by cataract and mild myotonia (sustained muscle contraction); life span is normal. Classic DM1 is characterized by muscle weakness and wasting, myotonia, cataract, and often by cardiac conduction abnormalities. Congenital DM1 is characterized by hypotonia and severe generalized weakness at birth, often with respiratory insufficiency and early death; intellectual disability is common. DM1 is caused by expansion of a CTG trinucleotide repeat in *DMPK*. Molecular genetic testing detects pathogenic variants in nearly 100% of affected individuals. DM1 is inherited in an autosomal dominant manner.
- **Myotonic dystrophy type 2 (DM2)**, also known as proximal myotonic myopathy (PROMM), is characterized by myotonia and muscle dysfunction (weakness, pain, and stiffness), and less commonly by cardiac conduction defects, iridescent posterior subcapsular cataracts, insulin-insensitive type 2 diabetes mellitus, and testicular failure. Although myotonia has been reported during the first decade, onset is typically in the third decade, most commonly with fluctuating or episodic muscle pain that can be debilitating and weakness of the neck flexors and finger flexors. Subsequently, weakness occurs in the elbow extensors and the hip flexors and extensors. Facial weakness and weakness of the ankle dorsiflexors are less common. Myotonia rarely causes severe symptoms. *CNBP* (*ZNF9*), the only gene known to be associated with DM2, has in intron 1 a complex repeat motif, (TG) $n$ (TCTG) $n$ (CCTG) $n$ , expansion of

which causes DM2. The number of CCTG repeats in expanded alleles ranges from approximately 75 to more than 11,000 (mean: ~5,000 repeats). The detection frequency of a *CNBP* CCTG expansion is more than 99%. DM2 is inherited in an autosomal dominant manner.

- **Motor and sensory neuropathies**
- **Inclusion body myopathy type 2 (IBM2)** is characterized by adult-onset, slowly progressive distal muscle weakness that begins with gait disturbance and foot drop secondary to anterior tibialis muscle weakness. Weakness eventually includes the hand and thigh muscles, but commonly spares the quadriceps muscles, even in advanced disease. Affected individuals are usually wheelchair bound about 20 years after onset. Muscle histopathology typically shows rimmed vacuoles and characteristic filamentous inclusions. *GNE*, which encodes the bifunctional enzyme UDP-N-acetylglucosamine 2-epimerase/N-acetylmannosamine kinase, is the only gene known to be associated with IBM2. IBM2 is inherited in an autosomal recessive manner.
- **Inclusion body myositis**, a slowly progressive and typically sporadic vacuolar myopathy, is associated with an autoaggressive inflammatory exudate and small congophilic inclusions.
- **Dysferlinopathy**, caused by pathogenic variants in *DYSF*, includes a spectrum of muscle disease characterized mainly by two phenotypes: Miyoshi myopathy, with primarily distal weakness; and limb-girdle muscular dystrophy type 2B (LGMD2B), with primarily proximal weakness. Miyoshi myopathy is characterized in young adults by muscle weakness and atrophy, most marked in the distal parts of the legs, especially the gastrocnemius and soleus muscles. Over a period of years, the weakness and atrophy spread to the thighs and gluteal muscles. LGMD2B is characterized by early weakness and atrophy of the pelvic and shoulder girdle muscles in adolescence or young adulthood, with slow progression. Dysferlinopathy is inherited in an autosomal recessive manner.
- **Other muscular dystrophies** that can predominantly affect distal muscles include [tibial muscular dystrophy](#) (Udd distal myopathy), telethoninopathy, [Laing distal myopathy](#) caused by *MYH7* pathogenic variants, Welander's distal dystrophy, [facioscapulohumeral dystrophy](#), anoctamin-5 related muscular dystrophy, and other sporadically occurring or dominantly inherited muscular dystrophies.

## Management

### Evaluations Following Initial Diagnosis

To establish the extent of disease in an individual diagnosed with myofibrillar myopathy (MFM), the following evaluations are recommended:

- EMG
- Routine ECG to identify arrhythmias and cardiac conduction defects; Holter monitoring if symptoms suggest an intermittent arrhythmia; echocardiogram if cardiac symptoms are present
- Respiratory function tests if respiratory symptoms are present
- Consultation with a clinical geneticist and/or genetic counselor

### Treatment of Manifestations

Pacemaker and implantable cardioverter defibrillator (ICD) should be considered in individuals with arrhythmia and/or cardiac conduction defects. Individuals with progressive or life-threatening cardiomyopathy are candidates for cardiac transplantation.

Respiratory support, consisting of continuous or bilevel positive airway pressure (CPAP or BIPAP), initially at night and later in the daytime, is indicated in individuals with hypercapnea and other signs of incipient ventilatory failure.



Range of motion physical therapy and assistive devices are appropriate for those with advanced muscle weakness. Treatment of scoliosis by spinal fusion is appropriate. Orthoses are indicated for treatment of foot drop.

## Prevention of Secondary Complications

Pacemaker or implantable cardioverter defibrillator placement for individuals with arrhythmogenic cardiomyopathy.

## Surveillance

The following are appropriate:

- Physical examination to monitor disease progression yearly or less often depending on rate of progression
- Electrocardiogram and/or echocardiogram yearly for early detection of cardiomyopathy
- Pulmonary function tests for individuals with exertional or nocturnal dyspnea.

## Evaluation of Relatives at Risk

Search [ClinicalTrials.gov](https://clinicaltrials.gov) in the US and [www.ClinicalTrialsRegister.eu](https://www.ClinicalTrialsRegister.eu) in Europe for information on clinical studies for a wide range of diseases and conditions. Note: There may not be clinical trials for this disorder.

## Pregnancy Management

Affected pregnant women may need assistance during delivery if they have significant weakness of their abdominal muscles.

## Therapies Under Investigation

Search [ClinicalTrials.gov](https://clinicaltrials.gov) in the US and [www.ClinicalTrialsRegister.eu](https://www.ClinicalTrialsRegister.eu) in Europe for access to information on clinical studies for a wide range of diseases and conditions. Note: There may not be clinical trials for this disorder.

## Other

The role of strengthening exercises has not been defined.

## Genetic Counseling

*Genetic counseling is the process of providing individuals and families with information on the nature, mode(s) of inheritance, and implications of genetic disorders to help them make informed medical and personal decisions. The following section deals with genetic risk assessment and the use of family history and genetic testing to clarify genetic status for family members; it is not meant to address all personal, cultural, or ethical issues that may arise or to substitute for consultation with a genetics professional. —ED.*

## Mode of Inheritance

Myofibrillar myopathy (MFM) is most commonly inherited in an autosomal dominant manner. Exceptions include:

- Autosomal recessive inheritance of *CRYAB* pathogenic frameshift variants that predict premature termination of the protein (see Molecular Genetics).
- X-linked inheritance of *FHL1* pathogenic variants

## Risk to Family Members – Autosomal Dominant Inheritance

### Parents of a proband

- Approximately 25% of individuals diagnosed with autosomal dominant (AD) myofibrillar myopathy have an affected parent.
- A proband with AD myofibrillar myopathy may have the disorder as the result of a *de novo* pathogenic variant. The proportion of cases caused by *de novo* pathogenic variants is unknown.
- Recommendations for the evaluation of parents of a proband with an apparent *de novo* pathogenic variant include clinical examination looking for weakness, EMG, and possible muscle biopsy. Molecular genetic testing may also be appropriate if a pathogenic variant has been identified in the proband.

Note: Twenty-five percent of individuals diagnosed with myofibrillar myopathy have an affected parent; in other individuals, the family history may appear to be negative because of failure to recognize the disorder in family members, early death of the parent before the onset of symptoms, or late onset of the disease in the affected parent.

### Sibs of a proband

- The risk to the sibs of the proband depends on the genetic status of the proband's parents.
- If a parent of the proband is affected, the risk to the sibs is 50%.
- When the parents are clinically unaffected, the risk to the sibs of a proband cannot be determined with confidence.
- Although no instances of germline mosaicism have been reported, it remains a possibility.

**Offspring of a proband.** Each child of an individual with AD myofibrillar myopathy has a 50% chance of inheriting the pathogenic variant.

**Other family members of a proband.** The risk to other family members depends on the status of the proband's parents: if a parent is affected or has a pathogenic variant, his or her family members are at risk.

## Risk to Family Members – Autosomal Recessive Inheritance

### Parents of a proband

- The parents of an affected individual are obligate heterozygotes (i.e., carriers of one mutated allele).
- Heterozygotes (carriers) are asymptomatic.

### Sibs of a proband

- At conception, each sib of an affected individual has a 25% chance of being affected, a 50% chance of being an asymptomatic carrier, and a 25% chance of being unaffected and not a carrier.
- Once an at-risk sib is known to be unaffected, the risk of his/her being a carrier is 2/3.
- Heterozygotes (carriers) are asymptomatic.

**Offspring of a proband.** Unless an individual with MFM has children with an affected individual or a carrier of a *CRYAB* pathogenic variant, his/her offspring will be obligate heterozygotes (carriers) for a pathogenic frameshift variant in *CRYAB*.

**Other family members.** Each sib of the proband's parents is at a 50% risk of being a carrier.

## Carrier (Heterozygote) Detection

Carrier testing for at-risk family members is possible if the *CRYAB* pathogenic variants in the family have been identified.

## Risk to Family Members – X-Linked Inheritance

### Parents of a proband

- The father of an affected male will not have MFM nor will he be a carrier of the *FHL1* pathogenic variant.
- In a family with more than one affected individual, the mother of an affected male is an obligate carrier.  
Note: If a woman has more than one affected child and no other affected relatives and if the pathogenic variant cannot be detected in her leukocyte DNA, she has germline mosaicism.
- If a male is the only affected family member (i.e., a simplex case), the mother may be a carrier or the affected male may have a *de novo* pathogenic variant and, thus, the mother is not a carrier. The percent of simplex cases representing *de novo* pathogenic variants is not known.

### Sibs of a proband

- The risk to sibs depends on the carrier status of the mother.
- If the mother of the proband has a *FHL1* pathogenic variant, the chance of transmitting it in each pregnancy is 50%. Males who inherit the pathogenic variant will be affected; females who inherit the pathogenic variant will be carriers and may be severely affected.
- If the proband represents a simplex case (i.e., a single occurrence in a family) and if the pathogenic variant cannot be detected in the leukocyte DNA of the mother, the risk to sibs is low but greater than that of the general population because of the possibility of maternal germline mosaicism.

**Offspring of a male proband.** Affected males pass the *FHL1* pathogenic variant to all of their daughters and none of their sons.

**Other family members.** The proband's maternal aunts may be at risk of being carriers and the aunts' offspring, depending on their gender, may be at risk of being carriers or of being affected.

Note: Molecular genetic testing may be able to identify the family member in whom a *de novo* pathogenic variant arose, information that could help determine genetic risk status of the extended family.

## Heterozygote (Carrier) Detection

Carrier testing for at-risk female relatives is possible if the *FHL1* pathogenic variant in the family has been identified.

## Related Genetic Counseling Issues

**Considerations in families with an apparent *de novo* pathogenic variant.** When neither parent of a proband with an autosomal dominant condition has clinical evidence of the disorder and/or the pathogenic variant, the proband likely has a *de novo* pathogenic variant. However, possible non-medical explanations including alternate paternity or maternity (e.g., with assisted reproduction) or undisclosed adoption could also be explored.

### Family planning

- The optimal time for determination of genetic risk, clarification of carrier status, and discussion of the availability of prenatal testing is before pregnancy.
- It is appropriate to offer genetic counseling (including discussion of potential risks to offspring and reproductive options) to young adults who are affected, are carriers, or are at risk of being affected or carriers.

**DNA banking** is the storage of DNA (typically extracted from white blood cells) for possible future use. Because it is likely that testing methodology and our understanding of genes, allelic variants, and diseases will improve in the future, consideration should be given to banking DNA of affected individuals.

## Prenatal Testing and Preimplantation Genetic Diagnosis

Once the pathogenic variant(s) have been identified in an affected family member, prenatal testing for a pregnancy at increased risk and preimplantation genetic diagnosis are possible.

## Resources

*GeneReviews staff has selected the following disease-specific and/or umbrella support organizations and/or registries for the benefit of individuals with this disorder and their families. GeneReviews is not responsible for the information provided by other organizations. For information on selection criteria, click [here](#).*

- Muscular Dystrophy Association - USA (MDA)**  
 222 South Riverside Plaza  
 Suite 1500  
 Chicago IL 60606  
**Phone:** 800-572-1717  
**Email:** [mda@mdausa.org](mailto:mda@mdausa.org)  
[www.mda.org](http://www.mda.org)
- Muscular Dystrophy UK**  
 61A Great Suffolk Street  
 London SE1 0BU  
 United Kingdom  
**Phone:** 0800 652 6352 (toll-free); 020 7803 4800  
**Email:** [info@muscular dystrophyuk.org](mailto:info@muscular dystrophyuk.org)  
[www.muscular dystrophyuk.org](http://www.muscular dystrophyuk.org)
- Sudden Arrhythmia Death Syndromes (SADS) Foundation**  
 508 East South Temple  
 Suite #202  
 Salt Lake City UT 84102  
**Phone:** 800-786-7723 (toll-free); 801-531-0937  
**Email:** [sads@sads.org](mailto:sads@sads.org)  
[www.sads.org](http://www.sads.org)

## Molecular Genetics

*Information in the Molecular Genetics and OMIM tables may differ from that elsewhere in the GeneReview: tables may contain more recent information.* —ED.

**Table A.** Myofibrillar Myopathy: Genes and Databases

Gene	Chromosome Locus	Protein	Locus-Specific Databases	HGMD	ClinVar
<a href="#">BAG3</a>	10q26.11	BAG family molecular chaperone regulator 3	<a href="#">BAG3 homepage - Leiden Muscular Dystrophy pages</a>	<a href="#">BAG3</a>	<a href="#">BAG3</a>
<a href="#">CRYAB</a>	11q23.1	Alpha-crystallin B chain	<a href="#">CRYAB homepage - Leiden Muscular Dystrophy pages</a>	<a href="#">CRYAB</a>	<a href="#">CRYAB</a>

Table A. continued from previous page.

<a href="#">DES</a>	<a href="#">2q35</a>	<a href="#">Desmin</a>	<a href="#">Human Intermediate Filament Database DES ARVD/C Genetic Variants Database (DES) DES homepage - Leiden Muscular Dystrophy pages</a>	<a href="#">DES</a>	<a href="#">DES</a>
<a href="#">DNAJB6</a>	<a href="#">7q36.3</a>	<a href="#">DnaJ homolog subfamily B member 6</a>	<a href="#">DNAJB6 homepage - Leiden Muscular Dystrophy pages</a>	<a href="#">DNAJB6</a>	<a href="#">DNAJB6</a>
<a href="#">FHL1</a>	<a href="#">Xq26.3</a>	<a href="#">Four and a half LIM domains protein 1</a>	<a href="#">FHL1 homepage - Leiden Muscular Dystrophy pages</a>	<a href="#">FHL1</a>	<a href="#">FHL1</a>
<a href="#">FLNC</a>	<a href="#">7q32.1</a>	<a href="#">Filamin-C</a>	<a href="#">FLNC homepage - Leiden Muscular Dystrophy pages</a>	<a href="#">FLNC</a>	<a href="#">FLNC</a>
<a href="#">LDB3</a>	<a href="#">10q23.2</a>	<a href="#">LIM domain-binding protein 3</a>	<a href="#">LDB3 homepage - Leiden Muscular Dystrophy pages</a>	<a href="#">LDB3</a>	<a href="#">LDB3</a>
<a href="#">MYOT</a>	<a href="#">5q31.2</a>	<a href="#">Myotilin</a>	<a href="#">MYOT homepage - Leiden Muscular Dystrophy pages</a>	<a href="#">MYOT</a>	<a href="#">MYOT</a>

Data are compiled from the following standard references: gene from [HGNC](#); chromosome locus from [OMIM](#); protein from [UniProt](#). For a description of databases (Locus Specific, HGMD, ClinVar) to which links are provided, click [here](#).

**Table B.** OMIM Entries for Myofibrillar Myopathy ([View All in OMIM](#))

<a href="#">102565</a>	<a href="#">FILAMIN C; FLNC</a>
<a href="#">123590</a>	<a href="#">CRYSTALLIN, ALPHA-B; CRYAB</a>
<a href="#">125660</a>	<a href="#">DESMIN; DES</a>
<a href="#">300163</a>	<a href="#">FOUR-AND-A-HALF LIM DOMAINS 1; FHL1</a>
<a href="#">601419</a>	<a href="#">MYOPATHY, MYOFIBRILLAR, 1; MFM1</a>
<a href="#">603883</a>	<a href="#">BCL2-ASSOCIATED ATHANOGENE 3; BAG3</a>
<a href="#">604103</a>	<a href="#">MYOTILIN; MYOT</a>
<a href="#">605906</a>	<a href="#">LIM DOMAIN-BINDING 3; LDB3</a>
<a href="#">608810</a>	<a href="#">MYOPATHY, MYOFIBRILLAR, 2; MFM2</a>
<a href="#">609200</a>	<a href="#">MYOPATHY, MYOFIBRILLAR, 3; MFM3</a>
<a href="#">609452</a>	<a href="#">MYOPATHY, MYOFIBRILLAR, 4; MFM4</a>
<a href="#">609524</a>	<a href="#">MYOPATHY, MYOFIBRILLAR, 5; MFM5</a>
<a href="#">611332</a>	<a href="#">DNAJ/HSP40 HOMOLOG, SUBFAMILY B, MEMBER 6; DNAJB6</a>

## Molecular Genetic Pathogenesis

Because myofibrillar myopathy is caused by pathogenic variants in any of eight different genes and each disease-associated gene may have different types of pathogenic variants, the molecular pathogenesis may vary from case to case. However, in all myofibrillar myopathies, the initial pathologic change involves disintegration of the Z-disk, and all disease proteins identified to date are involved in maintaining the structural integrity of the Z-disk. Because the Z-disks are sites of tension transmission between sarcomeres, the myofibrils fall apart when the Z-disks disintegrate.

**Animal models.** Desmin knockout mice show normal development of cardiac and skeletal muscle but subsequently show myofiber necrosis and phagocytosis [Capetanaki et al 1997]. This model is not comparable with human cases of desminopathy in which pathogenic missense variants weaken sarcomere structure and are associated with abnormal accumulation of desmin and other proteins. Transgenic mice that produce an inactivated form of human desmin (p.Arg173\_Glu179del) show desmin immunoreactive aggregates in myocardium [Muñoz-Mármol et al 1998]. Transgenic mice expressing an  $\alpha$ -B crystallin mutated protein (p.Arg120Gly) develop a severe cardiomyopathy with abnormal accumulation of desmin and  $\alpha$ -B crystallin in the heart; muscle pathology was not mentioned in this report [Wang et al 2001].

## DES

**Gene structure.** *DES* consists of nine exons. For a detailed summary of gene and protein information, see Table A, **Gene**.

**Pathogenic variants.** More than forty pathogenic variants including missense, frameshifting nucleotide insertion, small in-frame deletion, and splice-site variants have been reported [Goldfarb et al 2004]. See Table 2.

**Table 2.** Selected *DES* Pathogenic Variants

DNA Nucleotide Change	Predicted Protein Change	Reference Sequences
c.517_537del	p.Arg173_Glu179del	NM_001927.3 NP_001918.3

Note on variant classification: Variants listed in the table have been provided by the authors. *GeneReviews* staff have not independently verified the classification of variants.

Note on nomenclature: *GeneReviews* follows the standard naming conventions of the Human Genome Variation Society ([varnomen.hgvs.org](http://varnomen.hgvs.org)). See [Quick Reference](#) for an explanation of nomenclature.

**Normal gene product.** Desmin, comprising 470 amino acids, is a constituent of intermediate filaments in cardiac and skeletal muscle linking Z-disks with each other and to the subsarcolemmal cytoskeleton.

**Abnormal gene product.** Expression data suggest abnormal aggregation of mutated desmin molecules in heterologous systems. Pathogenic variants may interfere with desmin assembly and filament formation [Bär et al 2006].

## CRYAB

**Gene structure.** *CRYAB* consists of three exons. For a detailed summary of gene and protein information, see Table A, **Gene**.

**Pathogenic variants.** A nonsense variant, a missense variant, and a small frameshifting deletion have been reported [Vicart et al 1998, Selcen & Engel 2003]. Recently, autosomal recessive inheritance was shown for the p.Ser115ProfsTer14 [Forrest et al 2011] and Ser21AlafsTer24 [Del Bigio et al 2011] frameshift variants. See Table 3.



**Table 3.** Selected *CRYAB* Pathogenic Variants

DNA Nucleotide Change	Predicted Protein Change	Reference Sequences
c.60delC <sup>1</sup>	p.Ser21AlafsTer24	NM_001885.1 NP_001876.1
c.343delT <sup>2</sup>	p.Ser115ProfsTer14	
c.358A>G	p.Arg120Gly	

Note on variant classification: Variants listed in the table have been provided by the authors. *GeneReviews* staff have not independently verified the classification of variants.

Note on nomenclature: *GeneReviews* follows the standard naming conventions of the Human Genome Variation Society ([varnomen.hgvs.org](http://varnomen.hgvs.org)). See [Quick Reference](#) for an explanation of nomenclature.

1. Del Bigio et al [2011]

2. Homozygosity in an affected individual for this pathogenic variant in an autosomal recessive form of MFM [Forrest et al 2011]

**Normal gene product.** Alpha crystallin B chain ( $\alpha$ -B crystallin) is a small heat-shock chaperone protein required for maintaining the structural integrity of desmin. It is composed of 175 amino acid residues.

**Abnormal gene product.** Mutated alpha crystallin B chain molecules form smaller molecular-weight polymers than wild type in human muscle. In heterologous cells that constitutively express desmin, misfolded desmin molecules appear in aggregates.

## MYOT (TTID)

**Gene structure.** *MYOT* consists of ten exons. For a detailed summary of gene and protein information, see Table A, **Gene**.

**Pathogenic variants.** Five missense variants have been reported; all are in exon 2 [Hauser et al 2000, Hauser et al 2002, Selcen & Engel 2004]. See Table 4.

**Table 4.** Selected *MYOT* Pathogenic Variants

DNA Nucleotide Change	Predicted Protein Change	Reference Sequences
c.170C>T	p.Thr57Ile	NM_006790.2 NP_006781.1

Note on variant classification: Variants listed in the table have been provided by the authors. *GeneReviews* staff have not independently verified the classification of variants.

Note on nomenclature: *GeneReviews* follows the standard naming conventions of the Human Genome Variation Society ([varnomen.hgvs.org](http://varnomen.hgvs.org)). See [Quick Reference](#) for an explanation of nomenclature.

**Normal gene product.** Myotilin is a key Z-disk protein that interacts with  $\alpha$ -actinin, filamin-C, and actin. It has 498 amino acid residues.

**Abnormal gene product.** Mutated myotilin is predicted to weaken the linkage of Z-disk filaments to thin filaments. Transgenic mice with the p.Thr57Ile pathogenic variant and muscle disease similar to LGMD1A have been described [Garvey et al 2006].

## LDB3 (ZASP)

**Gene structure.** *LDB3* consists of 16 exons. It is alternatively spliced. For a detailed summary of gene and protein information, see Table A, **Gene**.

**Pathogenic variants.** Three missense variants have been reported [Selcen & Engel 2005]. See Table 5.

**Table 5.** Selected *LDB3* Pathogenic Variants

DNA Nucleotide Change	Predicted Protein Change	Reference Sequences
c.440C>T <sup>1</sup>	p.Ala147Val	<a href="#">NM_001080116.1</a> <a href="#">NP_001073585.1</a>

Note on variant classification: Variants listed in the table have been provided by the authors. *GeneReviews* staff have not independently verified the classification of variants.

Note on nomenclature: *GeneReviews* follows the standard naming conventions of the Human Genome Variation Society ([varnomen.hgvs.org](http://varnomen.hgvs.org)). See [Quick Reference](#) for an explanation of nomenclature.

1. Presents as a distal myopathy (see Genotype-Phenotype Correlations)

**Normal gene product.** LIM domain-binding protein 3 (Zasp) is a key Z-disk protein that interacts with  $\alpha$ -actinin and protein kinase C. It has 283 amino acid residues.

**Abnormal gene product.** Mutated LIM domain-binding protein 3 (Zasp) is predicted to weaken the linkage of Z-disk filaments to thin filaments.

## **FLNC**

**Gene structure.** Two *FLNC* transcript variants encoding different isoforms have been found for this gene (see Table A, **Gene**). The longer isoform, [NM\\_001458.4](#), consists of 48 exons.

**Pathogenic variants.** One nonsense variant and in-frame deletions have been reported [Vorgerd et al 2005, Shatunov et al 2009, Luan et al 2010].

**Normal gene product.** Filamin-C is a Z-disk protein that interacts with actin and myotilin. The protein [NP\\_001449.3](#) has 2725 amino acid residues.

**Abnormal gene product.** Mutated filamin-C has a disturbed secondary structure that prevents normal dimerization.

## **BAG3**

**Gene structure.** *BAG3* consists of four exons. No splice variants have been identified. For a detailed summary of gene and protein information, see Table A, **Gene**.

**Pathogenic variants.** The variant p.Pro209Leu has been reported [Selcen et al 2009]. See Table 6.

**Table 6.** Selected *BAG3* Pathogenic Variants

DNA Nucleotide Change	Predicted Protein Change	Reference Sequences
c.626C>T	p.Pro209Leu	<a href="#">NM_004281.3</a> <a href="#">NP_004272.2</a>

Note on variant classification: Variants listed in the table have been provided by the authors. *GeneReviews* staff have not independently verified the classification of variants.

Note on nomenclature: *GeneReviews* follows the standard naming conventions of the Human Genome Variation Society ([varnomen.hgvs.org](http://varnomen.hgvs.org)). See [Quick Reference](#) for an explanation of nomenclature.

**Normal gene product.** BAG family molecular chaperone regulator 3 (Bag3) is a co-chaperone for the Z-disk and has anti-apoptotic properties. It has 575 amino acid residues.

**Abnormal gene product.** Mutated Bag3 may alter the folding of Bag3 or may allosterically affect the binding properties of the canonical Bag3 domains.

## FHL1

**Gene structure.** *FHL1* has three alternatively spliced transcript variants that encode different protein isoforms with different tissue localizations and protein interactions. For a detailed summary of gene and protein information, see Table A, **Gene**.

**Pathogenic variants.** Selected reports of pathogenic variants include Quinzii et al [2008], Schessl et al [2008], Windpassinger et al [2008], Selcen et al [2011].

**Normal gene product.** FHL1 is a Z-disk related protein and some of its isoforms shuttle between cytoplasm and nucleus.

**Abnormal gene product.** Not known; likely dominant-negative effect.

## DNAJB6

**Gene structure.** *DNAJB6* has published two alternatively spliced transcript variants that encode different protein isoforms. The longer transcript variant, [NM\\_058246.3](#), has ten exons. For a detailed summary of gene and protein information, see Table A, **Gene**.

**Pathogenic variants.** Pathogenic variants have been reported [Harms et al 2012, Sarparanta et al 2012].

**Normal gene product.** The transcript [NM\\_005494.2](#) encodes a protein isoform with 241 amino acid residues ([NP\\_005485.1](#)). DnaJ homolog subfamily B member 6 (DNAJB6) is a member of heat shock protein family with molecular chaperone function.

**Abnormal gene product.** In vitro studies demonstrated that the pathogenic missense variants increase the half-life of DNAJB6 and reduce its anti-aggregation effect [Sarparanta et al 2012].

## References

### Literature Cited

- Abraham SC, DeNofrio D, Loh E, Minda JM, Tomaszewski JE, Pietra GG, Reynolds C. Desmin myopathy involving cardiac, skeletal, and vascular smooth muscle: report of a case with immunoelectron microscopy. *Hum Pathol*. 1998;29:876–82. PubMed PMID: 9712432.
- Arbustini E, Morbini P, Grasso M, Fasani R, Verga L, Bellini O, Dal Bello B, Campana C, Piccolo G, Febo O, Opasich C, Gavazzi A, Ferrans VJ. Restrictive cardiomyopathy, atrioventricular block and mild to subclinical myopathy in patients with desmin-immunoreactive material deposits. *J Am Coll Cardiol*. 1998;31:645–53. PubMed PMID: 9502648.
- Bär H, Mücke N, Ringler P, Müller SA, Kreplak L, Katus HA, Aebi U, Herrmann H. Impact of disease mutations on the desmin filament assembly process. *J Mol Biol*. 2006;360:1031–42. PubMed PMID: 16828798.
- Calderon A, Becker LE, Murphy EG. Subsarcolemmal vermiform deposits in skeletal muscle, associated with familial cardiomyopathy: report of two cases of a new entity. *Pediatr Neurosci*. 1987;13:108–12. PubMed PMID: 3438215.
- Capetanaki Y, Milner DJ, Weitzer G. Desmin in muscle formation and maintenance: knockouts and consequences. *Cell Struct Funct*. 1997;22:103–16. PubMed PMID: 9113396.
- Clark JR, D'Agostino AN, Wilson J, Brooks RR, Cole GC. Autosomal dominant myofibrillar inclusion body myopathy: clinical, histologic, histochemical, and ultrastructural characteristics. *Neurology*. 1978;28A:399.
- Dalakas MC, Park KY, Semino-Mora C, Lee HS, Sivakumar K, Goldfarb LG. Desmin myopathy, a skeletal myopathy with cardiomyopathy caused by mutations in the desmin gene. *N Engl J Med*. 2000;342:770–80. PubMed PMID: 10717012.

- De Bleecker JL, Engel AG, Ertl BB. Myofibrillar myopathy with abnormal foci of desmin positivity. II. Immunocytochemical analysis reveals accumulation of multiple other proteins. *J Neuropathol Exp Neurol*. 1996;55:563–77. PubMed PMID: 8627347.
- Del Bigio MR, Chudley AE, Sarnat HB, Campbell C, Goobie S, Chodirker BN, Selcen D. Infantile muscular dystrophy in Canadian aboriginals is an  $\alpha$ B-crystallinopathy. *Ann Neurol*. 2011;69:866–71. PubMed PMID: 21337604.
- Edström L, Thornell LE, Eriksson A. A new type of hereditary distal myopathy with characteristic sarcoplasmic bodies and intermediate (skeleton) filaments. *J Neurol Sci*. 1980;47:171–90. PubMed PMID: 6251174.
- Fardeau M, Godet-Guillain J, Tome FM, Collin H, Gaudeau S, Boffety C, Vernant P. *Rev Neurol (Paris)*. 1978;134:411–25. [A new familial muscular disorder demonstrated by the intra-sarcoplasmic accumulation of a granulo-filamentous material which is dense on electron microscopy (author's transl)]. PubMed PMID: 570292.
- Fidzianska A, Goebel HH, Osborn M, Lenard HG, Osse G, Langenbeck U. Mallory body-like inclusions in a hereditary congenital neuromuscular disease. *Muscle Nerve*. 1983;6:195–200. PubMed PMID: 6343859.
- Fischer D, Clemen CS, Olivé M, Ferrer I, Goudeau B, Roth U, Badorf P, Wattjes MP, Lutterbey G, Kral T, van der Ven PF, Fürst DO, Vicart P, Goldfarb LG, Moza M, Carpen O, Reichelt J, Schröder R. Different early pathogenesis in myotilinopathy compared to primary desminopathy. *Neuromuscul Disord*. 2006;16:361–7. PubMed PMID: 16684602.
- Forrest KM, Al-Sarraj S, Sewry C, Buk S, Tan SV, Pitt M, Durward A, McDougall M, Irving M, Hanna MG, Matthews E, Sarkozy A, Hudson J, Barresi R, Bushby K, Jungbluth H, Wraige E. Infantile onset myofibrillar myopathy due to recessive CRYAB mutations. *Neuromuscul Disord*. 2011;21:37–40. PubMed PMID: 21130652.
- Garvey SM, Miller SE, Claflin DR, Faulkner JA, Hauser MA. Transgenic mice expressing the myotilin T57I mutation unite the pathology associated with LGMD1A and MFM. *Hum Mol Genet*. 2006;15:2348–62. PubMed PMID: 16801328.
- Goebel HH, Muller J, Gillen HW, Merritt AD. Autosomal dominant "spheroid body myopathy". *Muscle Nerve*. 1978;1:14–26. PubMed PMID: 571956.
- Goebel HH. Desmin-related myopathies. *Curr Opin Neurol*. 1997;10:426–9. PubMed PMID: 9330890.
- Goldfarb LG, Park KY, Cervenáková L, Gorokhova S, Lee HS, Vasconcelos O, Nagle JW, Semino-Mora C, Sivakumar K, Dalakas MC. Missense mutations in desmin associated with familial cardiac and skeletal myopathy. *Nat Genet*. 1998;19:402–3. PubMed PMID: 9697706.
- Goldfarb LG, Vicart P, Goebel HH, Dalakas MC. Desmin myopathy. *Brain*. 2004;127:723–34. PubMed PMID: 14724127.
- Harms MB, Sommerville RB, Allred P, Bell S, Ma D, Cooper P, Lopate G, Pestronk A, Weihl CC, Baloh RH. Exome sequencing reveals DNAJB6 mutations in dominantly-inherited myopathy. *Ann Neurol*. 2012;71:407–16. PubMed PMID: 22334415.
- Hauser MA, Conde CB, Kowalchow V, Zeppa G, Taratuto AL, Torian UM, Vance J, Pericak-Vance MA, Speer MC, Rosa AL. myotilin Mutation found in second pedigree with LGMD1A. *Am J Hum Genet*. 2002;71:1428–32. PubMed PMID: 12428213.
- Hauser MA, Horrigan SK, Salmikangas P, Torian UM, Viles KD, Dancel R, Tim RW, Taivainen A, Bartoloni L, Gilchrist JM, Stajich JM, Gaskell PC, Gilbert JR, Vance JM, Pericak-Vance MA, Carpen O, Westbrook CA, Speer MC. Myotilin is mutated in limb girdle muscular dystrophy 1A. *Hum Mol Genet*. 2000;9:2141–7. PubMed PMID: 10958653.
- Horowitz SH, Schmalbruch H. Autosomal dominant distal myopathy with desmin storage: a clinicopathologic and electrophysiologic study of a large kinship. *Muscle Nerve*. 1994;17:151–60. PubMed PMID: 8114783.

- Kinoshita M, Satoyoshi E, Suzuki Y. Atypical myopathy with myofibrillar aggregates. *Arch Neurol*. 1975;32:417–20. PubMed PMID: 165803.
- Lobrinus JA, Janzer RC, Kuntzer T, Matthieu JM, Pfend G, Goy JJ, Bogousslavsky J. Familial cardiomyopathy and distal myopathy with abnormal desmin accumulation and migration. *Neuromuscul Disord*. 1998;8:77–86. PubMed PMID: 9608560.
- Luan X, Hong D, Zhang W, Wang Z, Yuan Y. A novel heterozygous deletion-insertion mutation (2695-2712 del/GTTTGT ins) in exon 18 of the filamin C gene causes filaminopathy in a large Chinese family. *Neuromuscul Disord*. 2010;20:390–6. PubMed PMID: 20417099.
- Muñoz-Mármol AM, Strasser G, Isamat M, Coulombe PA, Yang Y, Roca X, Vela E, Mate JL, Coll J, Fernández-Figueras MT, Navas-Palacios JJ, Ariza A, Fuchs E. A dysfunctional desmin mutation in a patient with severe generalized myopathy. *Proc Natl Acad Sci U S A*. 1998;95:11312–7. PubMed PMID: 9736733.
- Nakano S, Engel AG, Akiyuchi I, Kimura J. Myofibrillar myopathy. III. Abnormal expression of cyclin-dependent kinases and nuclear proteins. *J Neuropathol Exp Neurol*. 1997;56:850–6. PubMed PMID: 9258254.
- Nakashima N, Tamura Z, Okamoto S, Goto H. Inclusion bodies in human neuromuscular disorder. *Arch Neurol*. 1970;22:270–8. PubMed PMID: 5411681.
- Pénisson-Besnier I, Talvinen K, Dumez C, Vihola A, Dubas F, Fardeau M, Hackman P, Carpen O, Udd B. Myotilinopathy in a family with late onset myopathy. *Neuromuscul Disord*. 2006;16:427–31. PubMed PMID: 16793270.
- Quinzii CM, Vu TH, Min KC, Tanji K, Barral S, Grewal RP, Kattah A, Camaño P, Otaegui D, Kunitatsu T, Blake DM, Wilhelmsen KC, Rowland LP, Hays AP, Bonilla E, Hirano M. X-linked dominant scapuloperoneal myopathy is due to a mutation in the gene encoding four-and-a-half-LIM protein 1. *Am J Hum Genet*. 2008;82:208–13. PubMed PMID: 18179901.
- Sabatelli M, Bertini E, Ricci E, Salviati G, Magi S, Papacci M, Tonali P. Peripheral neuropathy with giant axons and cardiomyopathy associated with desmin type intermediate filaments in skeletal muscle. *J Neurol Sci*. 1992;109:1–10. PubMed PMID: 1517757.
- Sarparanta J, Jonson PH, Golzio C, Sandell S, Luque H, Screen M, McDonald K, Stajich JM, Mahjneh I, Vihola A, Raheem O, Penttilä S, Lehtinen S, Huovinen S, Palmio J, Tasca G, Ricci E, Hackman P, Hauser M, Katsanis N, Udd B. Mutations affecting the cytoplasmic functions of the co-chaperone DNAJB6 cause limb-girdle muscular dystrophy. *Nat Genet*. 2012;44:450–5. PubMed PMID: 22366786.
- Schessl J, Zou Y, McGrath MJ, Cowling BS, Maiti B, Chin SS, Sewry C, Battini R, Hu Y, Cottle DL, Rosenblatt M, Spruce L, Ganguly A, Kirschner J, Judkins AR, Golden JA, Goebel HH, Muntoni F, Flanigan KM, Mitchell CA, Bönnemann CG. Proteomic identification of FHL1 as the protein mutated in human reducing body myopathy. *J Clin Invest*. 2008;118:904–12. PubMed PMID: 18274675.
- Selcen D, Bromberg MB, Chin SS, Engel AG. Reducing bodies and myofibrillar myopathy features in FHL1 muscular dystrophy. *Neurology*. 2011;77:1951–9. PubMed PMID: 22094483.
- Selcen D, Engel AG. Mutations in myotilin cause myofibrillar myopathy. *Neurology*. 2004;62:1363–71. PubMed PMID: 15111675.
- Selcen D, Engel AG. Mutations in ZASP define a novel form of muscular dystrophy in humans. *Ann Neurol*. 2005;57:269–76. PubMed PMID: 15668942.
- Selcen D, Engel AG. Myofibrillar myopathy caused by novel dominant negative alpha B-crystallin mutations. *Ann Neurol*. 2003;54:804–10. PubMed PMID: 14681890.
- Selcen D, Muntoni F, Burton BK, Pegoraro E, Sewry C, Bite AV, Engel AG. Mutation in BAG3 causes severe dominant childhood muscular dystrophy. *Ann Neurol*. 2009;65:83–9. PubMed PMID: 19085932.

- Selcen D, Ohno K, Engel AG. Myofibrillar myopathy: clinical, morphological and genetic studies in 63 patients. *Brain*. 2004;127:439–51. PubMed PMID: 14711882.
- Shatunov A, Olivé M, Odgerel Z, Stadelmann-Nessler C, Irlbacher K, van Landeghem F, Bayarsaikhan M, Lee HS, Goudeau B, Chinnery PF, Straub V, Hilton-Jones D, Damian MS, Kaminska A, Vicart P, Bushby K, Dalakas MC, Sambuughin N, Ferrer I, Goebel HH, Goldfarb LG. In-frame deletion in the seventh immunoglobulin-like repeat of filamin C in a family with myofibrillar myopathy. *Eur J Hum Genet*. 2009;17:656–63. PubMed PMID: 19050726.
- Vicart P, Caron A, Guicheney P, Li Z, Prevost MC, Faure A, Chateau D, Chapon F, Tome F, Dupret JM, Paulin D, Fardeau M. A missense mutation in the alphaB-crystallin chaperone gene causes a desmin-related myopathy. *Nat Genet*. 1998;20:92–5. PubMed PMID: 9731540.
- Vorgerd M, van der Ven PF, Bruchertseifer V, Löwe T, Kley RA, Schröder R, Lochmüller H, Himmel M, Koehler K, Fürst DO, Huebner A. A mutation in the dimerization domain of filamin c causes a novel type of autosomal dominant myofibrillar myopathy. *Am J Hum Genet*. 2005;77:297–304. PubMed PMID: 15929027.
- Wang X, Osinska H, Klevitsky R, Gerdes AM, Nieman M, Lorenz J, Hewett T, Robbins J. Expression of R120G-alphaB-crystallin causes aberrant desmin and alphaB-crystallin aggregation and cardiomyopathy in mice. *Circ Res*. 2001;89:84–91. PubMed PMID: 11440982.
- Windpassinger C, Schoser B, Straub V, Hochmeister S, Noor A, Lohberger B, Farra N, Petek E, Schwarzbraun T, Ofner L, Löscher WN, Wagner K, Lochmüller H, Vincent JB, Quasthoff S. An X-linked myopathy with postural muscle atrophy and generalized hypertrophy, termed XMPMA, is caused by mutations in FHL1. *Am J Hum Genet*. 2008;82:88–99. PubMed PMID: 18179888.
- Wolburg H, Schlote W, Langohr HD, Peiffer J, Reiher KH, Heckl RW. Slowly progressive congenital myopathy with cytoplasmic bodies--report of two cases and a review of the literature. *Clin Neuropathol*. 1982;1:55–66. PubMed PMID: 6301720.

## Suggested Reading

- Olivé M. Extralysosomal protein degradation in myofibrillar myopathies. *Brain Pathol*. 2009;19:507–15. PubMed PMID: 19563542.
- Piñol-Ripoll G, Shatunov A, Cabello A, Larrodé P, de la Puerta I, Pelegrín J, Ramos FJ, Olivé M, Goldfarb LG. *Neuromuscul Disord*. 2009;19:418–22. PubMed PMID: 19433360.
- Schröder R, Schoser B. Myofibrillar myopathies: a clinical and myopathological guide. *Brain Pathol*. 2009;19:483–92. PubMed PMID: 19563540.
- Selcen D. Myofibrillar myopathies. *Curr Opin Neurol*. 2008;21:585–9. PubMed PMID: 18769253.
- Shalaby S, Mitsuhashi H, Matsuda C, Minami N, Noguchi S, Nonaka I, Nishino I, Hayashi YK. *J Neuropathol Exp Neurol*. 2009;68:701–7. PubMed PMID: 19458539.

## Chapter Notes

### Revision History

- 9 May 2019 (ma) Chapter retired: histologic diagnosis without strong genetic correlation
- 29 October 2012 (me) Comprehensive update posted live
- 27 July 2010 (cd) Revision: sequence analysis for *BAG3* mutations available clinically
- 2 February 2010 (me) Comprehensive update posted live
- 10 March 2008 (cd) Revision: sequence analysis and prenatal diagnosis available clinically for zaspopathy (*LDB3* mutations)
- 1 March 2007 (me) Comprehensive update posted live



- 9 January 2006 (ds) Revision: included disorders added (zaspopathy, filaminopathy)
- 1 July 2005 (ds) Revision: *DES* testing clinically available
- 28 January 2005 (me) Review posted live
- 2 August 2004 (ds) Original submission

## License

GeneReviews® chapters are owned by the University of Washington. Permission is hereby granted to reproduce, distribute, and translate copies of content materials for noncommercial research purposes only, provided that (i) credit for source (<http://www.genereviews.org/>) and copyright (© 1993-2024 University of Washington) are included with each copy; (ii) a link to the original material is provided whenever the material is published elsewhere on the Web; and (iii) reproducers, distributors, and/or translators comply with the [GeneReviews® Copyright Notice and Usage Disclaimer](#). No further modifications are allowed. For clarity, excerpts of GeneReviews chapters for use in lab reports and clinic notes are a permitted use.

For more information, see the [GeneReviews® Copyright Notice and Usage Disclaimer](#).

For questions regarding permissions or whether a specified use is allowed, contact: [admasst@uw.edu](mailto:admasst@uw.edu).