

Phenytoin Therapy and *HLA-B*15:02* and *CYP2C9* Genotype

Laura Dean, MD¹ and Megan Kane, PhD²

Created: September 22, 2016; Updated: April 7, 2021.

Introduction

Phenytoin (brand name Dilantin) is an anticonvulsant medication used for the treatment of seizures (1).

Phenytoin has a narrow therapeutic index—individuals that have supratherapeutic blood concentrations of phenytoin have increased risks of acute side effects. Dosing can be complex due to pharmacokinetic factors, including individual weight, age, gender, concomitant medications, plasma binding protein status, the presence of uremia or hyperbilirubinemia, and specific pharmacogenetic variants. As such, therapeutic drug monitoring is often used to adjust dose and maintain serum concentrations within the therapeutic range (10–20 µg/mL).

The *CYP2C9* enzyme is one of the main enzymes involved in the metabolism of phenytoin, and variant *CYP2C9* alleles are known to influence phenytoin drug levels. Individuals who have decreased activity *CYP2C9* variants may have reduced clearance rates of phenytoin and be at greater risk for dose-related side effects (2).


An individual's human leukocyte antigen B (*HLA-B*) genotype is a known risk factor for drug-induced hypersensitivity reactions. The *HLA-B* protein has an important immunological role in pathogen recognition and response, as well as to non-pathogens such as drugs. Individuals who have the *HLA-B*15:02* allele are at high risk of developing potentially life-threatening phenytoin-induced Stevens-Johnson syndrome (SJS) and the related toxic epidermal necrolysis (TEN).

The *HLA-B*15:02* allele is most often found among individuals of Southeast Asian descent, where there is a strong association between SJS/TEN and exposure to carbamazepine. Carbamazepine is an antiseizure medication used to treat the same types of seizures as phenytoin, as well as trigeminal neuralgia and bipolar disorder.

The FDA-approved drug label for phenytoin states that consideration should be given to avoiding phenytoin as an alternative for carbamazepine in individuals positive for *HLA-B*15:02* (Table 1). The label also mentions that variant *CYP2C9* alleles may contribute to unusually high levels of phenytoin (1).

Dosing recommendations for phenytoin based on *HLA-B* and *CYP2C9* genotype have also been published by the Clinical Pharmacogenetics Implementation Consortium (CPIC, Table 2, Figure 1) and the Dutch Pharmacogenetics Working Group (DPWG, Table 3, Table 4). These recommendations include the use of an antiseizure medication other than carbamazepine, phenytoin (or its prodrug fosphenytoin) for any *HLA-B*15:02* positive individual regardless of *CYP2C9* genotype, individual ancestry, or age. These recommendations also include specific dose reductions of phenytoin for individuals who have low or deficient enzyme activity (2, 3).

Clinical Pharmacogenetics Implementation Consortium (CPIC) Guideline for CYP2C9 and HLA-B Genotypes and Phenytoin Dosing: 2020 Update



Clinical Pharmacology & Therapeutics, First published: 11 August 2020, DOI: (10.1002/cpt.2008)

Figure 1: Dosage Guidelines from the CPIC for Phenytoin based on *HLA-B* and *CYP2C9* Genotype. Figure reproduced with permission from the authors.

Table 1. FDA Phenytoin Dosage and Administration (2019)

Gene or gene variant	Dosing considerations
<i>HLA-B*15:02</i>	Consideration should be given to avoiding phenytoin as an alternative for carbamazepine in individuals positive for <i>HLA-B*15:02</i> . The use of <i>HLA-B*15:02</i> genotyping has important limitations and must never substitute for appropriate clinical vigilance and individual management.
<i>CYP2C9</i> and <i>CYP2C19</i>	If individual is phenytoin naïve, do not use phenytoin/fosphenytoin. Avoid carbamazepine and oxcarbazepine. ^a If the individual has previously used phenytoin continuously for longer than 3 months without incidence of cutaneous adverse reactions, cautiously consider use of phenytoin in the future. The latency period for drug-induced SJS/TEN is short with continuous dosing and adherence to therapy (4–28 days), and cases usually occur within 3 months of dosing. ^b

This FDA table is adapted from (1).

Table 2. The CPIC Recommended Dosing of Phenytoin Based on *HLA-B*15:02* and *CYP2C9* Phenotype/Genotype (2020)

<i>CYP2C9</i> phenotype and <i>HLA-B</i> genotype	Implication	Therapeutic recommendation
Any <i>CYP2C9</i> phenotype and <i>HLA-B*15:02</i> positive [#]	Increased risk of phenytoin-induced SJS/ TEN	If the individual is phenytoin-naïve, do not use phenytoin/ fosphenytoin. Avoid carbamazepine and oxcarbazepine. ^a If the individual has previously used phenytoin continuously for longer than 3 months without incidence of cutaneous adverse reactions, cautiously consider use of phenytoin in the future. The latency period for drug-induced SJS/TEN is short with continuous dosing and adherence to therapy (428 days), and cases usually occur within 3 months of dosing. ^b
<i>CYP2C9</i> normal metabolizer and <i>HLA-B*15:02</i> negative	Normal phenytoin metabolism	No adjustments needed from typical dosing strategies. Subsequent doses should be adjusted according to therapeutic drug monitoring, response, and side effects. An <i>HLA-B*15:02</i> negative test does not eliminate the risk of phenytoin-induced SJS/TEN, and individuals should be carefully monitored according to standard practice. ^a
<i>CYP2C9</i> intermediate metabolizer (activity score 1.5) and <i>HLA-B*15:02</i> negative	Slightly reduced phenytoin metabolism: however, this does not appear to translate into increased side effects.	No adjustments needed from typical dosing strategies. Subsequent doses should be adjusted according to therapeutic drug monitoring, response, and side effects. An <i>HLA-B*15:02</i> negative test does not eliminate the risk of phenytoin-induced SJS/TEN, and individuals should be carefully monitored according to standard practice. ^c
<i>CYP2C9</i> intermediate metabolizer (Activity score 1.0) and <i>HLA-B*15:02</i> negative	Reduced phenytoin metabolism: Higher plasma concentrations will increase probability of toxicities.	For first dose, use typical initial or loading dose. For subsequent doses, use approximately 25% less than typical maintenance dose. Subsequent doses should be adjusted according to therapeutic drug monitoring, response, and side effects. An <i>HLA-B*15:02</i> negative test does not eliminate the risk of phenytoin-induced SJS/TEN, and individuals should be carefully monitored according to standard practice. ^c
<i>CYP2C9</i> poor metabolizer and <i>HLA-B*15:02</i> negative	Reduced phenytoin metabolism: Higher plasma concentrations will increase probability of toxicities.	For first dose, use typical initial or loading dose. For subsequent doses use approximately 50% less than typical maintenance dose. Subsequent doses should be adjusted according to therapeutic drug monitoring, response, and side effects. An <i>HLA-B*15:02</i> negative test does not eliminate the risk of phenytoin-induced SJS/TEN, and individuals should be carefully monitored according to standard practice. ^a

SJS/TEN: Stevens-Johnson syndrome/toxic epidermal necrolysis.

[#] Considerations: Other aromatic anticonvulsants, including eslicarbazepine, lamotrigine, and phenobarbital, have weaker evidence linking SJS/TEN with the *HLA-B*15:02* allele; however, caution should still be used in choosing an alternative agent. Previous tolerance of phenytoin is not indicative of tolerance to other aromatic anticonvulsants.

^a The strength of the therapeutic recommendation is classified as “strong”.

^b The strength of the therapeutic recommendation is classified as “optional”.

^c The strength of the therapeutic recommendation is classified as “moderate”.

This CPIC table is adapted from (2).

Note: The nomenclature used in this table reflects the standardized pharmacogenetic terms proposed by CPIC (4).

Table 3. The DPWG Phenytoin Dosing based on *HLA-B*15:02* Genotype (2017)

Genotype	Implication	Dosing recommendations
Positive for <i>HLA-B*15:02</i>	Increased risk of the life-threatening cutaneous side effect Stevens-Johnson syndrome/toxic epidermal necrolysis (SJS/TEN). The calculated risk of phenytoin-induced SJS/TEN in individuals with <i>HLA-B*15:02</i> is 0.65%.	<p>Carefully weigh the risk of SJS/TEN against the benefits Avoid phenytoin if an alternative is possible</p> <ul style="list-style-type: none"> • Carbamazepine has a 10-fold higher risk of SJS/TEN for these individuals and is therefore not an alternative. • A comparable risk has been reported for lamotrigine as for phenytoin. The same applies for oxcarbazepine, but the most severe forms (SJS/TEN overlap and TEN) are not observed with oxcarbazepine. <p>If it is not possible to avoid this medication, then advise the individual to report any skin rash immediately</p>

This DPWG table is adapted from (3).

Table 4. The DPWG Phenytoin Dosing based on *CYP2C9* Genotype (2018)

Metabolizer type	Genotype	Side effects	Recommendations		
			Loading dose	Other doses	Advise the individual
CYP2C9 IM	*1/*2	Genetic variation reduces conversion of phenytoin to inactive metabolites. This increases the risk of side effects. For the genotype group *1/*3+*3/*3, an increased risk of the life-threatening cutaneous side effects Stevens-Johnson Syndrome and toxic epidermal necrolysis has been observed, especially in Asians.	The loading dose does not need to be adjusted.	Use 75% of the standard dose*	Advise the individual to report side effects (such as ataxia, nystagmus, slurred speech, sedation or, especially in Asian individuals, rash) occur.
	*1/*3				
	other				
CYP2C9 PM	*2/*2			Use 50% of the standard dose*	
	*2/*3			Use 50% of the standard dose*	
	*3/*3			Use 40% of the standard dose*	
	other			Use 40–50% of the standard dose*	

* Assess the dose based on effect and serum concentration after 7–10 days.

This DPWG table is adapted from (5).

IM – intermediate metabolizer

PM – poor metabolizer

Drug: Phenytoin

Phenytoin is a generic antiseizure drug that is rarely prescribed to newly diagnosed individuals due to its propensity for long-term side effects. Nevertheless, it continues to be used by many individuals who initiated treatment before the availability of newer medications that have fewer side effects and drug-drug interactions. Phenytoin is used for the control of partial seizures and generalized tonic-clonic convulsions. It is also used in the treatment of status epilepticus and may be used to prevent or treat seizures that occur during and following neurosurgery (1).

Phenytoin belongs to the sodium channel blocker class of antiseizure drugs, which are thought to suppress seizure activity by blocking voltage-gated sodium channels that are responsible for the upstroke of action potentials (6, 7). The block by phenytoin and other members of this class of antiseizure drugs occurs in a state-dependent fashion, with preferential binding and block of the inactivated state of the channel. This results in

voltage- and frequency-dependent block in which high frequency action potential firing, which occurs during epileptic activity, is preferentially inhibited. (1, 8)

The dosing of phenytoin can be complex, as treatment is typically initiated at a low starting dose, which considers individual age, weight, and the presence of concomitant medications that may influence phenytoin metabolism or protein binding. The dose is then carefully escalated to obtain the desired therapeutic effect. There is a wide variation in how individuals respond to phenytoin (2). Therapeutic drug monitoring is often used to adjust the dose to ensure that plasma levels are within therapeutic range (10–20 µg/ml in adults). Measurement of plasma levels is useful when adding or discontinuing concomitant medications that effect phenytoin levels. Periodic measurement of plasma phenytoin concentrations may also be valuable in pregnancy because altered phenytoin pharmacokinetics increases the risk of seizures.

Phenytoin use during pregnancy has been associated with an 11% risk of fetal hydantoin syndrome in the offspring, which is characterized by dysmorphism, hypoplasia, and irregular ossification of the distal phalanges. Facial dysmorphism includes epicanthal folds, hypertelorism, broad flat nasal bridges, an upturned nasal tip, wide prominent lips, and, in addition, distal digital hypoplasia, intrauterine growth retardation, and mental retardation. An additional 30% of the in utero-exposed children express fetal hydantoin effects, in which there is a more limited pattern of dysmorphic characteristics. Some studies have found significant associations between in utero exposure to phenytoin and major congenital abnormalities (mainly, cardiac malformations and cleft palate) whereas others have failed to find such associations (9, 10).

The adverse effects of phenytoin fall into 2 categories, types A and B. (11)

Type A adverse drug reactions account for up to 90% of reactions. They are predictable and can occur in any individual if their drug exposure is high enough. Some of these reactions occur rapidly and are reversible when the dose is reduced. These include acute central nervous system adverse effects such as sedation, nystagmus, and ataxia. Other common side effects occur with long-term exposure and include changes to the physical appearance, such as gingival hyperplasia, coarsening of the facial features, hirsutism, and acne.

Type B adverse drug reactions include idiosyncratic hypersensitivity reactions, such as severe cutaneous adverse reaction (SCAR). Such reactions can occur at any dose and develop through a mechanism that is unrelated to the mechanism of action of the drug. A rare but life-threatening hypersensitivity reaction associated with phenytoin treatment is SJS and the related TEN. Both are severe cutaneous reactions to specific drugs, and are characterized by fever and lesions of the skin and mucous membranes, with a mortality rate of up to 30% (12).

It is difficult to predict in whom a drug-induced hypersensitivity reaction is likely to occur. However, for phenytoin individuals who are positive for a specific *HLA* allele are known to be susceptible to phenytoin-induced SJS/TEN. Human leukocyte antigen testing of individuals can identify at-risk individuals so that an alternative drug can be used.

The *HLA* Gene Family

The human leukocyte antigen (*HLA*) genes are members of the major histocompatibility complex (*MHC*) gene family, which includes more than 200 genes. The *MHC* family has been subdivided into 3 subgroups based on the structure and function of the encoded proteins: Class I, Class II, and Class III. The class I region contains the genes encoding the *HLA* molecules *HLA-A*, *HLA-B*, and *HLA-C*. These molecules are expressed on the surfaces of almost all cells and play an important role in processing and presenting antigens. The class I gene region also contains a variety of other genes, many of which are not known to be involved in immune function.

An important role of *HLA* class I molecules is to present peptide fragments to immune cells (CD8+ T cells). Most of these peptides originate from the breakdown of normal cellular proteins (“self”). However, if foreign peptide fragments are presented, for example, from a pathogen, CD8+T cells will recognize the peptides as “non-

self” and will be activated to release inflammatory cytokines and launch an immune response to dispose of the pathogen (or foreign body).

Because HLA molecules need to present such a wide variety of “self” and “non-self” peptides, the HLA genes are both numerous and highly polymorphic. More than 1,500 *HLA-B* alleles have been identified (13). The *HLA* allele nomenclature includes the HLA prefix, followed by the gene, an asterisk and a 2 digit number that corresponds to antigen specificity, and the assigned allele number (14). For example, the *HLA-B*15:02* allele is composed of:

- HLA: the HLA prefix (the HLA region on chromosome 6)
- B: the B gene (a particular HLA gene in this region)
- 15: the allele group (historically determined by serotyping, namely, a group of alleles that share the same serotype)
- 02: the specific HLA allele (a specific protein sequence; determined by genetic analysis).

Additional digits have recently been added to the nomenclature to discriminate alleles that do not differ in the protein amino acid sequence, but differ in their genetic sequence (namely, due to synonymous and noncoding genetic variants).

Variation in *HLA* genes plays an important role in the susceptibility to autoimmune disease and infections, and they are also critical in the context of transplant surgery where better outcomes are observed if the donor and recipient are HLA-compatible.

More recently, *HLA* alleles have been associated with susceptibility to Type B adverse drug reactions. For example, *HLA-B* alleles have been associated with severe hypersensitivity reactions to abacavir (used to treat HIV), allopurinol (used to treat gout), and the antiepileptic drugs carbamazepine and phenytoin.

Gene: *HLA-B*15:02*

Individuals who have one or 2 copies of the high-risk *HLA-B*15:02* allele are known as *HLA-B*15:02* positive (Table 5).

Table 5. The CPIC Assignment of Likely *HLA-B* Phenotype Based on Genotype (2020)

HLA phenotype	Genotype	Examples of diplotypes
<i>HLA-B*15:02</i> negative	Homozygous for any allele(s) other than <i>HLA-B*15:02</i>	*X/*X ^a
<i>HLA-B*15:02</i> positive	Homozygous or heterozygous variant	*15:02/*X ^a , *15:02/*15:02

^a Where *X = any *HLA-B* allele other than *15:02.

Table is adapted from (2).

Note: The nomenclature used in this table reflect the standardized pharmacogenetic terms proposed by CPIC (4).

The association between the *HLA-B*15:02* allele and SJS/TEN was first reported with the use of carbamazepine in the Han Chinese population. In the initial study, all individuals who had carbamazepine-induced SJS/TEN were found to have *HLA-B*15:02* (44/44, 100%), whereas the allele was much less common among carbamazepine-tolerant individuals (3/101, 3%)(15). In subsequent studies, this association was replicated, with an *HLA-B*15:02* positivity frequency of 70–100% among cases of carbamazepine-induced SJS/TEN (16).

The *HLA-B*15:02* allele was later associated with phenytoin-induced hypersensitivity reactions, including phenytoin-induced SJS in a Thai population and phenytoin-induced SJS/TEN in Chinese Asians (17, 18).

There are fewer studies on phenytoin-induced hypersensitivity than carbamazepine, and the strength of association between phenytoin and SJS/TEN is weaker than that of carbamazepine and SJS/TEN. However, from

the evidence available, the FDA recommends consideration of avoiding phenytoin as an alternative treatment to carbamazepine in individuals who have *HLA-B*15:02* (2).

The prevalence of carbamazepine-induced SJS/TEN is higher in populations where *HLA-B*15:02* is more common. Of note, the *HLA-B*15:02* allele frequency is highest in Southeast Asia, as populations from Hong Kong, Thailand, Malaysia, Vietnam, and parts of the Philippines have an allele frequency >15%. It is slightly lower (~10–13%) in Taiwan and Singapore, and around 4% in North China. South Asians, including Indians, appear to have an *HLA-B*15:02* allele frequency of ~2–4%, with higher frequencies in some subpopulations. (15, 16, 17, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30)

The *HLA-B*15:02* allele is rare (<1%) in East Asia (Japan and Korea) and among individuals who are not of Asian descent. For example, the allele is rare in Europeans, Hispanics, Africans, African Americans, and Native Americans. (16, 21)

Gene: **CYP2C9**

The cytochrome P450 superfamily (CYP450) is a large and diverse group of hepatic enzymes that form the major system for metabolizing lipids, hormones, toxins, and drugs. The CYP450 genes are very polymorphic and can result in reduced, absent, or increased enzyme activity.

The *CYP2C9* enzyme metabolizes approximately 15% of clinically used drugs, and atypical metabolic activity caused by genetic variants in the *CYP2C9* gene can play a major role in adverse drug reactions (31, 32).

The *CYP2C9* gene is polymorphic, with more than 50 known alleles. Variation in *CYP2C9* is thought to contribute to the pharmacogenetic variability in phenytoin metabolism. *CYP2C9*1* is considered the wild-type allele when no variants are detected and is categorized as normal enzyme activity (2). Individuals who have 2 normal-function alleles (for example, *CYP2C9*1/*1*) are classified as “normal metabolizers” (Table 6).

Table 6. The CPIC Assignment of Likely *CYP2C9* Phenotype Based on Genotype (2020)

<i>CYP2C9</i> phenotype ^{a,b}	Activity score	Genotype	Examples of diplotypes
Normal metabolizer	2	An individual with 2 normal-function alleles	*1/*1
Intermediate metabolizer	1.5 1	An individual with one normal-function allele plus one decreased-function allele OR one normal-function allele plus one no-function allele OR 2 decreased-function alleles	*1/*2 *1/*3 *2/*2
Poor metabolizer	0.5 0	An individual with one no-function allele plus one decreased-function allele; OR 2 no-function alleles	*2/*3 *3/*3

^a Assignment of allele function and associated citations can be found at the [CPIC website](#), also see (2).

^b See the *CYP2C9* Frequency Table in refs. 3 and 4 from (2) for population-specific allele and phenotype frequencies.

Note: There are no known cases of *CYP2C9* ultrarapid metabolizers

This CPIC table has been adapted from (2).

For individuals who are *CYP2C9* normal metabolizers, the recommended starting maintenance dose of phenytoin does not need to be adjusted based on genotype (2).

Two common allelic variants associated with reduced enzyme activity are *CYP2C9*2* (p.Arg144Cys) and *CYP2C9*3* (p.Ile359Leu). The *2 allele is more common in Caucasian (10–20%), than Asian (1–3%) or African (0–6%) populations. The *3 allele is less common (<10% in most populations) and is extremely rare in African populations. In African Americans, the *CYP2C9*5*, *6, *8 and *11 alleles are more common (33, 34, 35).

Linking *HLA-B* and *CYP2C9* Genetic Variation with the Risk of Side Effects and Treatment Response

Reduced activity *CYP2C9* alleles, in particular *CYP2C9**3, influence phenytoin dosage, individual response, and predict adverse drug reactions (36, 37, 38, 39, 40). Individuals with reduced-function *CYP2C9* alleles have reduced clearance of phenytoin and have an increased risk of side effects.

Specific HLA alleles, namely, *HLA-B**15:02, are also strongly associated with SCAR (41).

To guide the optimal dose of phenytoin and reduce the risk, both genetic factors (*CYP2C9* and *HLA* alleles) and non-genetic factors (for example, omeprazole co-medication) need to be considered (41, 42, 43, 44, 45).

The *HLA-B* and *CYP2C9* Gene Interactions with Medications Used for Additional Indications

Other medications with multiple indications are known to interact with the *HLA-B* alleles or to be metabolized by *CYP2C9*.

- Other seizure medications—Carbamazepine has similar interactions with HLA variation and hypersensitivity reactions: individuals with one or more copies of *HLA-B**15:02 are at risk of SJS/TEN and the *HLA-A**31:01 is strongly associated with a potentially life-threatening condition known as drug reaction with eosinophilia and systemic symptoms (DRESS) and a milder reaction maculopapular exanthema (MPE).
- Uric acid reduction medications—Allopurinol is a xanthine oxidase inhibitor to treat high uric acid levels seen in gout, tumor lysis syndrome, and cases of symptomatic hyperuricemia. The *HLA-B**58:01 allele is associated with SCAR during allopurinol treatment. Lesinurad is a urate transport inhibitor, also used in the treatment of gout and it is metabolized by *CYP2C9* to inactive metabolites. Individuals who are *CYP2C9* poor metabolizers will have an increased exposure to the active drug and thus have an increased risk of side effects such as kidney stones or cardiovascular events.
- Anti-retroviral medication—Abacavir, a nucleoside/nucleotide reverse transcriptase inhibitor used in the treatment of HIV, is also associated with hypersensitivity reactions, the risk of which increases when an individual has the *HLA-B**57:01 allele.
- Non-steroidal anti-inflammatory drugs (NSAIDs)—Celecoxib, Flurbiprofen, and Piroxicam are used for pain management in osteoarthritis, rheumatoid arthritis, and other conditions; they are all metabolized by *CYP2C9* and poor metabolizers will experience higher levels of exposure to these NSAIDs and have higher risk of side effects.
- Anti-emetics—Dronabinol, a synthetic cannabinoid, is used in the treatment of chemotherapy-induced nausea and vomiting for individuals who had poor responses to traditional anti-emetics. It also is used to treat anorexia-associated weight loss in individuals with acquired immunodeficiency syndrome. Dronabinol is activated by *CYP2C9* metabolism and individuals with poor metabolizer phenotypes will have an increased exposure to dronabinol and increased risk of side effects.

Additional information on gene-drug interactions for *HLA-B* and *CYP2C9* are available from [PharmGKB](#), [CPIC](#), and the [FDA](#) (search for “*HLA-B*” or “*CYP2C9*”).

Genetic Testing

The NIH’s Genetic Testing Registry provides examples of the genetic tests that are available for the [phenytoin](#) drug response, the [HLA-B](#) gene, and the [CYP2C9](#) gene.

The genotype results for an *HLA* allele such as *HLA-B*15:02* can either be “positive” or “negative”. There are no intermediate phenotypes because the *HLA* genes are expressed in a codominant manner.

A positive result indicates the individual is either “heterozygous” or “homozygous” for the variant, depending upon whether they have one or 2 copies of the **15:02* allele.

A negative result indicates that the individual does not have the *HLA-B*15:02* allele. However, a negative result does not rule out the possibility of an individual developing phenytoin-induced SJS/TEN. Therefore, clinicians should carefully monitor all individuals according to standard practices.

For *CYP2C9*, alleles to be included in clinical genotyping assays have been recommended by the Association for Molecular Pathology. (46) Results are typically reported as a diplotype, such as *CYP2C9 *1/*2*, and may include an interpretation of the individual’s predicted metabolizer phenotype (normal, intermediate, or poor) and an activity score (Table 6).

Therapeutic Recommendations based on Genotype

This section contains excerpted¹ information on gene-based dosing recommendations. Neither this section nor other parts of this review contain the complete recommendations from the sources.

2019 Statement from the US Food and Drug Administration (FDA)

Regarding *HLA-B*:

Studies in individuals of Chinese ancestry have found a strong association between the risk of developing SJS/TEN and the presence of *HLA-B*15:02*, an inherited allelic variant of the *HLA-B* gene, in individuals using carbamazepine. Limited evidence suggests that *HLA-B*15:02* may be a risk factor for the development of SJS/TEN in individuals of Asian ancestry taking other antiepileptic drugs associated with SJS/TEN, including phenytoin. Consideration should be given to avoiding phenytoin as an alternative for carbamazepine in individuals positive for *HLA-B*15:02*.

The use of *HLA-B*15:02* genotyping has important limitations and must never substitute for appropriate clinical vigilance and individual management. The role of other possible factors in the development of, and morbidity from, SJS/TEN, such as antiepileptic drug (AED) dose, compliance, concomitant medications, comorbidities, and the level of dermatologic monitoring have not been studied.

Regarding *CYP2C9* and *CYP2C19*:

In most individuals maintained at a steady dosage, stable phenytoin serum levels are achieved. There may be wide interindividual variability in phenytoin serum levels with equivalent dosages. Individuals with unusually low levels may be noncompliant or hypermetabolizers of phenytoin.

Unusually high levels result from liver disease, variant *CYP2C9* and *CYP2C19* alleles, or drug interactions which result in metabolic interference. The individual with large variations in phenytoin serum levels, despite standard doses, presents a difficult clinical problem. Serum level determinations in such individuals may be particularly helpful. As phenytoin is highly protein bound, free phenytoin levels may be altered in individuals whose protein binding characteristics differ from normal.

Please review the complete therapeutic recommendations that are located here: (1).

¹ The FDA labels specific drug formulations. We have substituted the generic names for any drug labels in this excerpt. The FDA may not have labeled all formulations containing the generic drug.

2018 Summary of recommendations from the Dutch Pharmacogenetics Working Group (DPWG) of the Royal Dutch Association for the Advancement of Pharmacy (KNMP)

HLA-B*1502

The life-threatening cutaneous side effect Stevens-Johnson syndrome/toxic epidermal necrolysis (SJS/TEN) occurs more frequently in patients with this genetic variation. The calculated risk of phenytoin-induced SJS/TEN in patients with *HLA-B*15:02* is 0.65%.

- Carefully weigh the risk of SJS/TEN against the benefits
- Avoid phenytoin if an alternative is possible
 - Carbamazepine carries a 10-fold higher risk of SJS/TEN for these individuals and is therefore not an alternative.
 - A comparable risk has been reported for lamotrigine as for phenytoin. The same applies for oxcarbazepine, but the most severe forms (SJS/TEN overlap and TEN) are not observed with oxcarbazepine.
- If it is not possible to avoid this medication, then advise the individual to report any skin rash immediately (Table 2)

CYP2C9 genotypes *1/*2, *1/*3 and other IMs

Genetic variation reduces conversion of phenytoin to inactive metabolites. This increases the risk of side effects.

Recommendation:

1. The loading dose does not need to be adjusted.
2. For the other doses, use 75% of the standard dose and assess the dose based on effect and serum concentration after 7-10 days.
3. Advise the patient to report if side effects (such as ataxia, nystagmus, slurred speech, sedation or rash) occur.

CYP2C9 genotypes *2/*2 and *2/*3 and other PMs

Genetic variation reduces conversion of phenytoin to inactive metabolites. This increases the risk of side effects.

Recommendation:

1. The loading dose does not need to be adjusted.
2. For the other doses, use 50% of the standard dose and assess the dose based on effect and serum concentration after 7-10 days.
3. Advise the patient to report if side effects (such as ataxia, nystagmus, slurred speech, sedation or rash) occur.

CYP2C9 genotype *3/*3 (PM)

Genetic variation reduces conversion of phenytoin to inactive metabolites. This increases the risk of side effects, including SJS/TEN in Asian patients.

Recommendation:

1. The loading dose does not need to be adjusted.
2. For the other doses, use 40% of the standard dose and assess the dose based on effect and serum concentration after 7-10 days.

3. Advise the patient to report if side effects (such as ataxia, nystagmus, slurred speech, sedation or, especially in Asian individuals, rash) occur.

Please review the complete therapeutic recommendations that are located here (3).

2020 Statement from the Clinical Pharmacogenetics Implementation Consortium (CPIC)

*HLA-B*15:02* recommendations

[...] If a individual is phenytoin-naïve and *HLA-B*15:02* positive, the individual has an increased risk of SJS/TEN and the recommendation is to consider using an anticonvulsant other than phenytoin unless the benefits of treating the underlying disease clearly outweigh the risks (see Table 3). Carbamazepine and oxcarbazepine should also be avoided if a individual is *HLA-B*15:02* positive. Alternative medications such as eslicarbazepine acetate and lamotrigine have limited evidence linking SJS/TEN with the *HLA-B*15:02* allele.

[...]

If a individual is phenytoin-naïve and *HLA-B*15:02* negative, the individual has a normal risk of phenytoin-induced SJS/TEN and the recommendation is to use phenytoin with dosage adjustments based on *CYP2C9* genotype (if known) or standard dosing guidelines (if *CYP2C9* genotype is unknown). However, an *HLA-B*15:02* negative test does not eliminate the risk of phenytoin-induced SJS/TEN.

CYP2C9 recommendations.

The recommended phenytoin initial or loading and maintenance doses do not need adjustments based on genotype for *CYP2C9* NMs and IMs with an AS of ≥ 1.5 . Available evidence does not clearly indicate the extent of dose reduction needed to prevent phenytoin-related toxicities in *CYP2C9* IMs with an AS of 1.0 and PMs with an AS of 0 or 0.5. Furthermore, multiple case studies have observed that *CYP2C9* PMs are at increased risk for exposure-related phenytoin toxicities, and multiple studies have observed an association between the *CYP2C9*3* allele and SJS/TEN. Although presence of the *CYP2C9*3* allele is insufficient to predict phenytoin-induced SJS/TEN, these and other data suggest that the risk of SJS/TEN is dose-related and provide an additional rationale for reducing phenytoin dose in *CYP2C9* PMs. Thus, our recommendations are conservative given the variability surrounding phenytoin dosing. Based on the doses reported in the pharmacokinetic and pharmacogenetic studies mentioned above and in Table S2, a typical initial or loading dose followed by at least a 25% reduction in the recommended starting maintenance dose may be considered for *CYP2C9* IMs with AS of 1.0. Subsequent maintenance doses should be adjusted based on therapeutic drug monitoring and response. For *CYP2C9* PMs, use a typical initial or loading dose then consider at least a 50% reduction of starting maintenance dose with subsequent maintenance doses adjusted based on therapeutic drug monitoring and response.

Pediatrics

Much of the evidence (summarized in Table S1) linking *HLA-B*15:02* to phenytoin induced SJS/TEN was generated in both children and adults. Therefore, the above recommendation is made regardless of *CYP2C9* genotype, individual age, race or ancestry.

Please review the complete therapeutic recommendations that are located here: (2).

Nomenclature of Selected *HLA-B* Alleles

Allele name	dbSNP reference identifier for allele location
<i>HLA-B*15:02</i>	Tagged variants cannot be reliably used to detect this allele. Sequencing is the most accurate technique for allele detection.

For the *major histocompatibility complex* region, variations in genes such as *HLA-B* occur across the whole sequence of the gene, not a single locus. Therefore, the *HLA-B*15:02* allele is defined by its sequence rather than single coding or protein variations. If there is strong linkage disequilibrium between one or more SNPs and a specific *HLA* allele, the presence of these SNPs (tag SNPs) may be used for *HLA* typing in some populations; however, genotyping tagged variants should not be considered diagnostic or equivalent to actual *HLA* testing. For *HLA-B*15:02*, rs2844682 and rs3909184 were previously considered the tagged variants (47), however, these variants have shown to be less accurate in other studies (48). Other tagged variants have been suggested, however, the sensitivity and accuracy of these variants to detect the *HLA-B*15:02* allele is limited (49, 50). Sequence for the full *HLA-B*15:02* allele (and subtypes) can be accessed [here](#).

Guidelines on nomenclature of the *HLA* system are available from *HLA* Nomenclature: <http://hla.alleles.org/>

Nomenclature of Selected *CYP2C9* Alleles

Common allele name	Alternative names	HGVS reference sequence		dbSNP reference identifier for allele location
		Coding	Protein	
<i>CYP2C9*2</i>	430C>T Arg144Cys	NM_000771.3:c.430C>T	NP_000762.2:p.Arg144Cys	rs1799853
<i>CYP2C9*3</i>	1075A>C Ile359Leu	NM_000771.3:c.1075A>C	NP_000762.2:p.Ile359Leu	rs1057910
<i>CYP2C9*5</i>	1080C>G Asp360Glu	NM_000771.3:c.1080C>G	NP_000762.2:p.Asp360Glu	rs28371686
<i>CYP2C9*6</i>	817delA Lys273Argfs	NM_000771.3:c.818del	NP_000762.2:p.Lys273Argfs	rs9332131
<i>CYP2C9*8</i>	449G>A Arg150His	NM_000771.3:c.449G>A	NP_000762.2:p.Arg150His	rs7900194
<i>CYP2C9*11</i>	1003C>T Arg335Trp	NM_000771.3:c.1003C>T	NP_000762.2:p.Arg335Trp	rs28371685

Guidelines for the description and nomenclature of gene variations are available from the Human Genome Variation Society (HGVS): <http://www.hgvs.org/content/guidelines>

Nomenclature for Cytochrome P450 enzymes is available from the Pharmacogene Variation Consortium (PharmVar) <https://www.pharmvar.org/>.

Acknowledgments

The authors would like to thank Marga Nijenhuis, PhD, Royal Dutch Pharmacists Association (KNMP), The Hague, The Netherlands; Bernard Esquivel MD, PhD, President of the Latin American Association for Personalized Medicine, Vancouver, BC, Canada; and Jason H. Karnes, PharmD, PhD, BCPS, FAHA, Assistant Professor, Department of Pharmacy Practice & Science, University of Arizona College of Pharmacy; Sarver Heart Center, University of Arizona College of Medicine, Tucson, Arizona, USA for reviewing this summary

First edition:

The author would like to thank Mohamed Nagy, Clinical Pharmacist, Head of the Personalised Medication Management Unit, Department of Pharmaceutical Services, Children's Cancer Hospital, Cairo, Egypt; Emily K. Pauli, Director of Research, Clearview Cancer Institute, Huntsville, AL, USA; Michael A. Rogawski, Professor of

Neurology, University of California, Davis, CA, USA; and Stuart A. Scott, Assistant Professor of Genetics and Genomic Sciences, Icahn School of Medicine at Mount Sinai, New York, NY, USA for reviewing this summary.

Version History

To view the previous version of this chapter, published on 22 September 2016, please click [here](#).

References

1. PHENYTOIN suspension [package insert]. Morton Grove, IL: Morton Grove Pharmaceuticals Inc; 2019. Available from: <https://dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?setid=efd93f07-818b-41ae-abd6-49ec5175311a>
2. Karnes J.H., Rettie A.E., Somogyi A.A., Huddart R., et al. Clinical Pharmacogenetics Implementation Consortium (CPIC) Guideline for CYP2C9 and HLA-B Genotypes and Phenytoin Dosing: 2020 Update. *Clin Pharmacol Ther.* 2021;109(2):302–309. PubMed PMID: 32779747.
3. Royal Dutch Pharmacists Association (KNMP). Dutch Pharmacogenetics Working Group (DPWG). Pharmacogenetic Guidelines [Internet]. Netherlands. Phenytoin – HLA-B*1502 [Cited July 2020]. Available from: <http://kennisbank.knmp.nl>
4. Caudle K.E., Dunnenberger H.M., Freimuth R.R., Peterson J.F., et al. Standardizing terms for clinical pharmacogenetic test results: consensus terms from the Clinical Pharmacogenetics Implementation Consortium (CPIC). *Genet Med.* 2017;19(2):215–223. PubMed PMID: 27441996.
5. Royal Dutch Pharmacists Association (KNMP). Dutch Pharmacogenetics Working Group (DPWG). Pharmacogenetic Guidelines [Internet]. Netherlands. Phenytoin – CYP2C9 [Cited July 2020]. Available from: <http://kennisbank.knmp.nl>
6. Catterall W.A. Molecular properties of brain sodium channels: an important target for anticonvulsant drugs. *Adv Neurol.* 1999;79:441–56. PubMed PMID: 10514834.
7. Lipkind G.M., Fozzard H.A. Molecular model of anticonvulsant drug binding to the voltage-gated sodium channel inner pore. *Mol Pharmacol.* 2010;78(4):631–8. PubMed PMID: 20643904.
8. Segal M.M., Douglas A.F. Late sodium channel openings underlying epileptiform activity are preferentially diminished by the anticonvulsant phenytoin. *J Neurophysiol.* 1997;77(6):3021–34. PubMed PMID: 9212254.
9. *Birth Defects: Data & Statistics*. Centers for Disease Control and Prevention (CDC); Available from: <http://www.cdc.gov/ncbddd/birthdefects/data.html>.
10. Hill D.S., Wlodarczyk B.J., Palacios A.M., Finnell R.H. Teratogenic effects of antiepileptic drugs. *Expert Rev Neurother.* 2010;10(6):943–59. PubMed PMID: 20518610.
11. Mullan K.A., Anderson A., Illing P.T., Kwan P., et al. HLA-associated antiepileptic drug-induced cutaneous adverse reactions. *HLA.* 2019;93(6):417–435. PubMed PMID: 30895730.
12. Svensson C.K., Cowen E.W., Gaspari A.A. Cutaneous drug reactions. *Pharmacol Rev.* 2001;53(3):357–79. PubMed PMID: 11546834.
13. Nomenclature for Factors of the HLA System: HLA Alleles [Cited 23 June 2016]. Available from: <http://hla.alleles.org/alleles/index.html>
14. Choo S.Y. The HLA system: genetics, immunology, clinical testing, and clinical implications. *Yonsei Med J.* 2007;48(1):11–23. PubMed PMID: 17326240.
15. Chung W.H., Hung S.I., Hong H.S., Hsieh M.S., et al. Medical genetics: a marker for Stevens-Johnson syndrome. *Nature.* 2004;428(6982):486. PubMed PMID: 15057820.
16. Amstutz U., Shear N.H., Rieder M.J., Hwang S., et al. Recommendations for HLA-B*15:02 and HLA-A*31:01 genetic testing to reduce the risk of carbamazepine-induced hypersensitivity reactions. *Epilepsia.* 2014;55(4):496–506. PubMed PMID: 24597466.
17. Locharearnkul C., Loplumert J., Limotai C., Korkij W., et al. Carbamazepine and phenytoin induced Stevens-Johnson syndrome is associated with HLA-B*1502 allele in Thai population. *Epilepsia.* 2008;49(12):2087–91. PubMed PMID: 18637831.

18. Hung S.I., Chung W.H., Liu Z.S., Chen C.H., et al. Common risk allele in aromatic antiepileptic-drug induced Stevens-Johnson syndrome and toxic epidermal necrolysis in Han Chinese. *Pharmacogenomics*. 2010;11(3):349–56. PubMed PMID: 20235791.
19. TEGRETOL (carbamazepine) tablet [package insert]. East Hanover, New Jersey 07936: Corporation, N.P.; 2011. Available from: <http://dailymed.nlm.nih.gov/dailymed/lookup.cfm?setid=8d409411-aa9f-4f3a-a52c-fbcb0c3ec053>
20. Leckband S.G., Kelsoe J.R., Dunnenberger H.M., George A.L. Jr, et al. Clinical Pharmacogenetics Implementation Consortium guidelines for HLA-B genotype and carbamazepine dosing. *Clin Pharmacol Ther*. 2013;94(3):324–8. PubMed PMID: 23695185.
21. Chung W.H., Hung S.I., Chen Y.T. Genetic predisposition of life-threatening antiepileptic-induced skin reactions. *Expert Opin Drug Saf*. 2010;9(1):15–21. PubMed PMID: 20001755.
22. Puangpetch A., Koomdee N., Chamnanphol M., Jantararoungtong T., et al. HLA-B allele and haplotype diversity among Thai patients identified by PCR-SSOP: evidence for high risk of drug-induced hypersensitivity. *Front Genet*. 2014;5:478. PubMed PMID: 25657656.
23. Nguyen D.V., Chu H.C., Nguyen D.V., Phan M.H., et al. HLA-B*1502 and carbamazepine-induced severe cutaneous adverse drug reactions in Vietnamese. *Asia Pac Allergy*. 2015;5(2):68–77. PubMed PMID: 25938071.
24. Chong K.W., Chan D.W., Cheung Y.B., Ching L.K., et al. Association of carbamazepine-induced severe cutaneous drug reactions and HLA-B*1502 allele status, and dose and treatment duration in paediatric neurology patients in Singapore. *Arch Dis Child*. 2014;99(6):581–4. PubMed PMID: 24225276.
25. Hung S.I., Chung W.H., Jee S.H., Chen W.C., et al. Genetic susceptibility to carbamazepine-induced cutaneous adverse drug reactions. *Pharmacogenet Genomics*. 2006;16(4):297–306. PubMed PMID: 16538176.
26. Lonjou C., Thomas L., Borot N., Ledger N., et al. A marker for Stevens-Johnson syndrome ...: ethnicity matters. *Pharmacogenomics J*. 2006;6(4):265–8. PubMed PMID: 16415921.
27. Alfirevic A., Jorgensen A.L., Williamson P.R., Chadwick D.W., et al. HLA-B locus in Caucasian patients with carbamazepine hypersensitivity. *Pharmacogenomics*. 2006;7(6):813–8. PubMed PMID: 16981842.
28. Kaniwa N., Saito Y., Aihara M., Matsunaga K., et al. HLA-B locus in Japanese patients with anti-epileptics and allopurinol-related Stevens-Johnson syndrome and toxic epidermal necrolysis. *Pharmacogenomics*. 2008;9(11):1617–22. PubMed PMID: 19018717.
29. Mehta T.Y., Prajapati L.M., Mittal B., Joshi C.G., et al. Association of HLA-B*1502 allele and carbamazepine-induced Stevens-Johnson syndrome among Indians. *Indian J Dermatol Venereol Leprol*. 2009;75(6):579–82. PubMed PMID: 19915237.
30. Wu X.T., Hu F.Y., An D.M., Yan B., et al. Association between carbamazepine-induced cutaneous adverse drug reactions and the HLA-B*1502 allele among patients in central China. *Epilepsy Behav*. 2010;19(3):405–8. PubMed PMID: 20833111.
31. Van Booven D., Marsh S., McLeod H., Carrillo M.W., et al. Cytochrome P450 2C9-CYP2C9. *Pharmacogenet Genomics*. 2010;20(4):277–81. PubMed PMID: 20150829.
32. Gupta A., Zheng L., Ramanujam V., Gallagher J. Novel Use of Pharmacogenetic Testing in the Identification of CYP2C9 Polymorphisms Related to NSAID-Induced Gastropathy. *Pain Med*. 2015;16(5):866–9. PubMed PMID: 25585969.
33. Sistonen J., Fuselli S., Palo J.U., Chauhan N., et al. Pharmacogenetic variation at CYP2C9, CYP2C19, and CYP2D6 at global and microgeographic scales. *Pharmacogenet Genomics*. 2009;19(2):170–9. PubMed PMID: 19151603.
34. Solus J.F., Arietta B.J., Harris J.R., Sexton D.P., et al. Genetic variation in eleven phase I drug metabolism genes in an ethnically diverse population. *Pharmacogenomics*. 2004;5(7):895–931. PubMed PMID: 15469410.
35. Lee C.R., Goldstein J.A., Pieper J.A. Cytochrome P450 2C9 polymorphisms: a comprehensive review of the in-vitro and human data. *Pharmacogenetics*. 2002;12(3):251–63. PubMed PMID: 11927841.

36. Liao K., Liu Y., Ai C.Z., Yu X., et al. The association between CYP2C9/2C19 polymorphisms and phenytoin maintenance doses in Asian epileptic patients: A systematic review and meta-analysis. *Int J Clin Pharmacol Ther.* 2018;56(7):337–346. PubMed PMID: 29628024.
37. Fohner A.E., Ranatunga D.K., Thai K.K., Lawson B.L., et al. Assessing the clinical impact of CYP2C9 pharmacogenetic variation on phenytoin prescribing practice and patient response in an integrated health system. *Pharmacogenet Genomics.* 2019;29(8):192–199. PubMed PMID: 31461080.
38. Calderon-Ospina C.A., Galvez J.M., Lopez-Cabra C., Morales N., et al. Possible Genetic Determinants of Response to Phenytoin in a Group of Colombian Patients With Epilepsy. *Front Pharmacol.* 2020;11:555. PubMed PMID: 32457604.
39. Wen Y.F., Culhane-Pera K.A., Thyagarajan B., Bishop J.R., et al. Potential Clinical Relevance of Differences in Allele Frequencies Found within Very Important Pharmacogenes between Hmong and East Asian Populations. *Pharmacotherapy.* 2020;40(2):142–152. PubMed PMID: 31884695.
40. Fohner A.E., Rettie A.E., Thai K.K., Ranatunga D.K., et al. Associations of CYP2C9 and CYP2C19 Pharmacogenetic Variation with Phenytoin-Induced Cutaneous Adverse Drug Reactions. *Clin Transl Sci.* 2020;13(5):1004–1009. PubMed PMID: 32216088.
41. Chang W.C., Hung S.I., Carleton B.C., Chung W.H. An update on CYP2C9 polymorphisms and phenytoin metabolism: implications for adverse effects. *Expert Opin Drug Metab Toxicol.* 2020;16(8):723–734. PubMed PMID: 32510242.
42. Yampayon K., Sukasem C., Limwongse C., Chinvarun Y., et al. Influence of genetic and non-genetic factors on phenytoin-induced severe cutaneous adverse drug reactions. *Eur J Clin Pharmacol.* 2017;73(7):855–865. PubMed PMID: 28391407.
43. Su S.C., Chen C.B., Chang W.C., Wang C.W., et al. HLA Alleles and CYP2C9*3 as Predictors of Phenytoin Hypersensitivity in East Asians. *Clin Pharmacol Ther.* 2019;105(2):476–485. PubMed PMID: 30270535.
44. Hikino K., Ozeki T., Koido M., Terao C., et al. HLA-B*51:01 and CYP2C9*3 Are Risk Factors for Phenytoin-Induced Eruption in the Japanese Population: Analysis of Data From the Biobank Japan Project. *Clin Pharmacol Ther.* 2020;107(5):1170–1178. PubMed PMID: 31646624.
45. Sukasem C., Sririttha S., Tempark T., Klaewsongkram J., et al. Genetic and clinical risk factors associated with phenytoin-induced cutaneous adverse drug reactions in Thai population. *Pharmacoepidemiol Drug Saf.* 2020;29(5):565–574. PubMed PMID: 32134161.
46. Pratt V.M., Cavallari L.H., Del Tredici A.L., Hachad H., et al. Recommendations for Clinical CYP2C9 Genotyping Allele Selection: A Joint Recommendation of the Association for Molecular Pathology and College of American Pathologists. *J Mol Diagn.* 2019;21(5):746–755. PubMed PMID: 31075510.
47. de Bakker P.I., McVean G., Sabeti P.C., Miretti M.M., et al. A high-resolution HLA and SNP haplotype map for disease association studies in the extended human MHC. *Nat Genet.* 2006;38(10):1166–72. PubMed PMID: 16998491.
48. Zhu G.D., Brenton A.A., Malhotra A., Riley B.J., et al. Genotypes at rs2844682 and rs3909184 have no clinical value in identifying HLA-B*15:02 carriers. *Eur J Clin Pharmacol.* 2015;71(8):1021–3. PubMed PMID: 26036218.
49. He Y., Hoskins J.M., Clark S., Campbell N.H., et al. Accuracy of SNPs to predict risk of HLA alleles associated with drug-induced hypersensitivity events across racial groups. *Pharmacogenomics.* 2015;16(8):817–24. PubMed PMID: 26083016.
50. Fang H., Xu X., Kaur K., Dedek M., et al. A Screening Test for HLA-B (*)15:02 in a Large United States Patient Cohort Identifies Broader Risk of Carbamazepine-Induced Adverse Events. *Front Pharmacol.* 2019;10:149. PubMed PMID: 30971914.

License

All Medical Genetics Summaries content, except where otherwise noted, is licensed under a Creative Commons Attribution 4.0 International (CC BY 4.0) license which permits copying, distribution, and adaptation of the work,

provided the original work is properly cited and any changes from the original work are properly indicated. Any altered, transformed, or adapted form of the work may only be distributed under the same or similar license to this one.