




Icaridin

Revised: December 3, 2018.

CASRN: 119515-38-7



Drug Levels and Effects

Summary of Use during Lactation

No information is available on the clinical use of icaridin (picaridin) during breastfeeding. However, the Centers for Disease Control and Prevention and U.S. Environmental Protection Agency consider icaridin to be safe and effective during breastfeeding when used as directed. It should be used by breastfeeding women to avoid exposure to mosquito-borne viruses.[1] Avoid application directly to the nipple and other areas where the infant might directly ingest the product.

Drug Levels

Maternal Levels. Relevant published information was not found as of the revision date.

Disclaimer: Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site.

Attribution Statement: LactMed is a registered trademark of the U.S. Department of Health and Human Services.

Infant Levels. Relevant published information was not found as of the revision date.

Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

Alternate Drugs to Consider

Citriodiol, Diethyltoluamide, Insect Repellent M 3535

References

1. Centers for Disease Control and Prevention. Avoid bug bites. Accessed 6-27/16. Available at: <http://wwwnc.cdc.gov/travel/page/avoid-bug-bites#repellent>

Substance Identification

Substance Name

Icaridin

CAS Registry Number

119515-38-7

Drug Class

Breast Feeding

Lactation

Insect Repellents