

respectively. The authors estimated that a fully breastfed would receive a dose of propafenone and its metabolite of about 0.03% of the mother's weight-adjusted dosage.[1]

One mother (time postpartum not stated) took a single dose of propafenone 150 mg orally. The highest milk propafenone level was 37.4 mcg/L at 2 hours after the dose. By 6 hours after the dose, the drug was undetectable (<10 mcg/L). The highest milk 5-hydroxypropafenone level was 102 mcg/L at 2 hours after the dose. By 6 hours after the dose, the level was 19.8 mcg/L and by 12 hours it was undetectable (<10 mcg/L) in milk. The authors estimated that an exclusively breastfed infant would receive 0.1% of the maternal weight-adjusted dosage of propafenone as drug and metabolite.[2]

Infant Levels. Relevant published information was not found as of the revision date.

Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

References

1. Libardoni M, Piovan D, Busato E et al. Transfer of propafenone and 5-OH-propafenone to foetal plasma and maternal milk. *Br J Clin Pharmacol.* 1991;32:527-8. PubMed PMID: 1958453.
2. Wakaumi M, Tsuruoka S, Sakamoto K et al. Pilsicainide in breast milk from a mother: comparison with disopyramide and propafenone. *Br J Clin Pharmacol.* 2005;59:120-2. PubMed PMID: 15606453.

Substance Identification

Substance Name

Propafenone

CAS Registry Number

54063-53-5

Drug Class

Breast Feeding

Lactation

Antiarrhythmics