



Varicella Vaccine

Revised: February 15, 2024.

Drug Levels and Effects

Summary of Use during Lactation

The Centers for Disease Control and Prevention and the several health professional organizations state that vaccines given to a nursing mother do not affect the safety of breastfeeding for mothers or infants and that breastfeeding is not a contraindication to varicella vaccine. Breastfed infants should be vaccinated according to the routine recommended schedules.[1-3]

Drug Levels

Maternal Levels. Two postpartum women immunized with varicella vaccine provided preimmunization and serial postimmunization breastmilk samples (number not specified). One sample of colostrum contained detectable viral DNA and B-actin by polymerase chain reaction, but all other samples were negative for these species. No varicella gene sequences were found in any of the samples.[4]

One study of 12 women vaccinated with live, attenuated varicella vaccine found no evidence of varicella virus excretion into breastmilk.[5]

Infant Levels. Relevant published information was not found as of the revision date.

Effects in Breastfed Infants

Limited data indicate that breastfeeding can enhance the response of the infant to certain vaccine antigens. [1,2,6]

A 25-day-old otherwise healthy infant developed a rash that was found to be caused by the varicella vaccine strain. The baby's mother received the varicella vaccine (Varivax, Merck & Co) on day 3 postpartum, when the infant was present in the room. Neither the mother nor any known contact had a rash prior to the infant's rash. No other known contact had been vaccinated in the 90 days prior to the infant's presentation at the clinic. The infant was primarily formula-fed, but received expressed breastmilk 4 times a day. The authors considered transmission via breastmilk, saliva or respiratory spread unlikely, but possible. The most likely exposure was thought to be via aerosolization at the time of vaccine administration.[7]

Disclaimer: Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site.

Attribution Statement: LactMed is a registered trademark of the U.S. Department of Health and Human Services.

Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

References

1. Ezeanolue E, Harriman K, Hunter P, et al. Best Practices Guidance of the Advisory Committee on Immunization Practices (ACIP). Available at: <https://www.cdc.gov/vaccines/hcp/acip-recs/general-recs/index.html>
2. Kimberlin DW, Brady MT, Jackson MA, Long SS. Red Book: 2018 Report of the Committee on Infectious Diseases. 31st ed. Elk Grove Village, IL: American Academy of Pediatrics. 2018.
3. Gruslin A, Steben M, Halperin S, et al. Immunization in pregnancy: No. 220, December 2008. *Int J Gynaecol Obstet* 2009;105:187-91. PubMed PMID: 19367691.
4. Dolbear GL, Moffat J, Falkner C, Wojtowycz M. A pilot study: Is attenuated varicella virus present in breast milk after postpartum immunization? *Obstet Gynecol* 2003;101 (4 Suppl):47s. Abstract.
5. Bohlke K, Galil K, Jackson LA, et al. Postpartum varicella vaccination: is the vaccine virus excreted in breast milk? *Obstet Gynecol* 2003;102:970-7. PubMed PMID: 14672472.
6. Pabst HF. Immunomodulation by breast-feeding. *Pediatr Infect Dis J* 1997;16:991-5. PubMed PMID: 9380478.
7. Kluthe M, Herrera A, Blanca H, et al. Neonatal vaccine-strain varicella-zoster virus infection 22 days after maternal postpartum vaccination. *Pediatr Infect Dis J* 2012;31:977-9. PubMed PMID: 22572750.

Substance Identification

Substance Name

Varicella Vaccine

Drug Class

Breast Feeding

Lactation

Milk, Human

Vaccines