




## Acetaminophen

Revised: February 15, 2024.

CASRN: 103-90-2



## Drug Levels and Effects

### Summary of Use during Lactation

Acetaminophen is a good choice for analgesia, and fever reduction in nursing mothers. Amounts in milk are much less than doses usually given to infants. Adverse effects in breastfed infants appear to be rare.

### Drug Levels

*Maternal Levels.* A single oral dose of 650 mg of acetaminophen was given to 12 nursing mothers who were 2 to 22 months postpartum. Peak milk levels of 10 to 15 mg/L occurred between 1 and 2 hours after the dose in all patients. Acetaminophen was undetectable (<0.5 mg/L) in all mothers 12 hours after the dose. The authors calculated that an infant who ingested 90 mL of breastmilk every 3 hours would receive an average of 0.88 mg of

**Disclaimer:** Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site.

**Attribution Statement:** LactMed is a registered trademark of the U.S. Department of Health and Human Services.

acetaminophen or 0.14% (range 0.04 to 0.23%) of the mother's absolute dosage.[1] Using data from this study, an infant would receive a maximum of about 2% of the maternal weight-adjusted dosage.

Three women took a single 500 mg dose of acetaminophen. Peak milk levels averaging 4.2 mg/L occurred within 2 hours after the dose.[2] Using data from this study, an infant would receive a maximum of about 3.6% of the maternal weight-adjusted dosage.

Four women who were 2 to 8 months postpartum were given a single 1 gram dose of acetaminophen. Milk was completely emptied from one breast every 30 minutes for 3 to 3.5 hours, with a final sample from the opposite breast. Peak milk levels occurred between 1 and 2.5 hours after the dose. The acetaminophen level in the breast that was sampled only once had a lower level than the breast sampled at half-hour intervals. The authors estimated that a breastfed infant would receive an average of 1.1% and a maximum of 1.8% of the maternal weight-adjusted dosage. This dose is about 0.5% of the lowest recommended infant dose of acetaminophen.[3]

Eighteen women taking acetaminophen 500 mg orally postpartum pain donated a total of 20 milk samples for analysis at random times at 5 to 6 days postpartum. Milk concentrations ranged from 2.5 mcg/L at 68 hours after a dose to 13.5 mg/L at 0.8 hours after a dose. Levels were generally lower with increased time after a dose.[4] Note that this paper does not state the number of doses taken prior to sampling, so steady-state was not guaranteed. The pharmacokinetic analysis of samples was performed in an invalid way, and pharmacokinetic parameters reported in the paper are therefore not valid.

*Infant Levels.* No acetaminophen was detected in the urine of 12 breastfed infants aged 2 to 22 months after maternal ingestion of 650 mg of acetaminophen.[1]

Urine was collected for 1 to 3.5 hours after nursing in 6 infants aged 2 to 6 days whose mothers received 1 to 2 grams of acetaminophen 2 to 4 hours before nursing their infant. Infants excreted an average of 401 mcg of acetaminophen and its metabolites in urine during the collection interval. These neonates excreted a greater percentage of drug as acetaminophen and much less as the sulfate metabolite than in an adult comparison population.[3]

## Effects in Breastfed Infants

A maculopapular rash on the upper trunk and face of a 2-month-old infant was probably caused by acetaminophen in breastmilk. The rash occurred after 2 days of therapy in the mother at a dose of 1 gram at bedtime. It subsided when the drug was discontinued and recurred 2 weeks later after another acetaminophen dose of 1 gram was taken by the mother.[5]

Two papers report 14 women who breastfed after taking acetaminophen or its prodrug phenacetin with no adverse effects to their infants.[6,7]

In a telephone follow-up study, mothers reported no side effects among 43 infants exposed to acetaminophen in breastmilk.[8]

Two clinicians speculated that breastmilk exposure to acetaminophen during breastfeeding might be a risk factor for asthma and wheezing in the breastfed infants based on their personal observations.[9] However, these observations were uncontrolled and cannot be considered to be valid proof of an association.[10]

## Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

## Alternate Drugs to Consider

Ibuprofen

## References

1. Berlin CM, Jr, Yaffe SJ, Ragni M. Disposition of acetaminophen in milk, saliva and plasma of lactating women. *Pediatr Pharmacol (New York)* 1980;1:135-41. PubMed PMID: 7202185.
2. Bitzén PO, Gustafsson B, Jostell KG, et al. Excretion of paracetamol in human breast milk. *Eur J Clin Pharmacol* 1981;20:123-5. PubMed PMID: 7262173.
3. Notarianni LJ, Oldham HG, Bennett PN. Passage of paracetamol into breast milk and its subsequent metabolism by the neonate. *Br J Clin Pharmacol* 1987;24:63-7. PubMed PMID: 3620287.
4. Tamaki R, Noshiro K, Furugen A, et al. Breast milk concentrations of acetaminophen and diclofenac - unexpectedly high mammary transfer of the general-purpose drug acetaminophen. *BMC Pregnancy Childbirth* 2024;24:90. PubMed PMID: 38287321.
5. Matheson I, Lunde PKM, Notarianni L. Infant rash caused by paracetamol in breast milk? *Pediatrics* 1985;76:651-2. PubMed PMID: 2931668.
6. Berlin CM, Jr, Yaffe SJ, Ragni M. Disposition of acetaminophen in milk, saliva and plasma of lactating women. *Pediatr Pharmacol* 1980;1:135-41. PubMed PMID: 7202185.
7. Findlay JW, DeAngelis RL, Kearney MF, et al. Analgesic drugs in breast milk and plasma. *Clin Pharmacol Ther* 1981;29:625-33. PubMed PMID: 7214793.
8. Ito S, Blajchman A, Stephenson M, et al. Prospective follow-up of adverse reactions in breast-fed infants exposed to maternal medication. *Am J Obstet Gynecol* 1993;168:1393-9. PubMed PMID: 8498418.
9. Verd S, Nadal-Amat J. Paracetamol and asthma and lactation. *Acta Paediatr* 2011;100:e2-3. PubMed PMID: 21244486.
10. Bakkeheim E, Carlsen KH, Lodrup Carlsen, KC. Paracetamol exposure during breastfeeding and risk of allergic disease. *Acta Paediatr* 2011;100:e3. doi:10.1111/j.1651-2227.2011.02340.x

## Substance Identification

### Substance Name

Acetaminophen

### CAS Registry Number

103-90-2

### Drug Class

Breast Feeding

Lactation

Milk, Human

Analgesics, Non-Narcotic