




Cefditoren

Revised: August 15, 2023.

CASRN: 104145-95-1



Drug Levels and Effects

Summary of Use during Lactation

Cefditoren is no longer marketed in the United States. Although no information is available on the use of cefditoren during breastfeeding, cephalosporins are generally not expected to cause adverse effects in breastfed infants. Occasionally disruption of the infant's gastrointestinal flora, resulting in diarrhea or thrush have been reported with cephalosporins, but these effects have not been adequately evaluated. Cefditoren is acceptable in nursing mothers.

Drug Levels

Maternal Levels. Relevant published information was not found as of the revision date.

Disclaimer: Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site.

Attribution Statement: LactMed is a registered trademark of the U.S. Department of Health and Human Services.

Infant Levels. Relevant published information was not found as of the revision date.

Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

Alternate Drugs to Consider

Ceftibuten

Substance Identification

Substance Name

Cefditoren

CAS Registry Number

104145-95-1

Drug Class

Breast Feeding

Lactation

Milk, Human

Anti-Infective Agents

Antibacterial Agents

Cephalosporins