




Cromolyn

Revised: April 15, 2023.

CASRN: 15826-37-6



Drug Levels and Effects

Summary of Use during Lactation

Although no published data exist on the use of cromolyn during lactation, maternal milk levels are likely to be very low and it is expected to be poorly absorbed from the infant's gastrointestinal tract. An expert panel considers use of cromolyn to be acceptable during breastfeeding.[1]

Drug Levels

Maternal Levels. Relevant published information was not found as of the revision date.

Infant Levels. Relevant published information was not found as of the revision date.

Disclaimer: Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site.

Attribution Statement: LactMed is a registered trademark of the U.S. Department of Health and Human Services.

Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

References

1. National Heart, Lung, and Blood, Institute, et al. NAEPF expert panel report. Managing asthma during pregnancy: recommendations for pharmacologic treatment-2004 update. 2004;1-57.

Substance Identification

Substance Name

Cromolyn

CAS Registry Number

15826-37-6

Drug Class

Breast Feeding

Lactation

Milk, Human

Anti-Allergic Agents

Anti-Asthmatic Agents