




Lodoxamide

Revised: December 3, 2018.

CASRN: 53882-12-5



Drug Levels and Effects

Summary of Use during Lactation

Although no published data exist on the use of lodoxamide during lactation, maternal milk levels are likely to be very low after the use eye drops. To substantially diminish the amount of drug that reaches the breastmilk after using eye drops, place pressure over the tear duct by the corner of the eye for 1 minute or more, then remove the excess solution with an absorbent tissue.

Drug Levels

Maternal Levels. Relevant published information was not found as of the revision date.

Infant Levels. Relevant published information was not found as of the revision date.

Disclaimer: Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site.

Attribution Statement: LactMed is a registered trademark of the U.S. Department of Health and Human Services.

Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

Substance Identification

Substance Name

Lodoxamide

CAS Registry Number

53882-12-5

Drug Class

Breast Feeding

Lactation

Anti-Allergic Agents

Histamine Antagonists