




## Fidaxomicin

Revised: May 17, 2021.

CASRN: 873857-62-6



## Drug Levels and Effects

### Summary of Use during Lactation

No information is available on the use of fidaxomicin during breastfeeding. Because it is poorly absorbed orally, it is not likely to reach the bloodstream of the infant or cause any adverse effects in breastfed infants.

### Drug Levels

*Maternal Levels.* Relevant published information was not found as of the revision date.

*Infant Levels.* Relevant published information was not found as of the revision date.

**Disclaimer:** Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site.

**Attribution Statement:** LactMed is a registered trademark of the U.S. Department of Health and Human Services.

## Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

## Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

## Alternate Drugs to Consider

Vancomycin

## Substance Identification

### Substance Name

Fidaxomicin

### CAS Registry Number

873857-62-6

### Drug Class

Breast Feeding

Lactation

Anti-Infective Agents

Macrocyclic Antibiotics