




## Plazomicin

Revised: August 16, 2021.

CASRN: 1154757-24-0



## Drug Levels and Effects

### Summary of Use during Lactation

Plazomicin is an aminoglycoside antibiotic similar to gentamicin and amikacin. No information is available on the use of plazomicin during breastfeeding. However, based on the excretion of other aminoglycoside antibiotics, amounts in milk are expected to be low. Monitor the infant for possible effects on the gastrointestinal flora, such as diarrhea, candidiasis (e.g., thrush, diaper rash) or rarely, blood in the stool indicating possible antibiotic-associated colitis.

### Drug Levels

*Maternal Levels.* Relevant published information was not found as of the revision date.

**Disclaimer:** Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site.

**Attribution Statement:** LactMed is a registered trademark of the U.S. Department of Health and Human Services.

*Infant Levels.* Relevant published information was not found as of the revision date.

### **Effects in Breastfed Infants**

Relevant published information was not found as of the revision date.

### **Effects on Lactation and Breastmilk**

Relevant published information was not found as of the revision date.

## **Substance Identification**

### **Substance Name**

Plazomicin

### **CAS Registry Number**

1154757-24-0

### **Drug Class**

Breast Feeding

Lactation

Antibacterial Agents

Anti-Infective Agents

Aminoglycosides