



Monoclonal Antibodies

Updated: February 7, 2024.

OVERVIEW

Monoclonal antibodies are immunoglobulins that have a high degree of specificity (mono-specificity) for an antigen or epitope. Monoclonal antibodies are typically derived from a clonal expansion of antibody producing malignant human plasma cells. The initial monoclonal antibodies were created by fusing spleen cells from an immunized mouse with human or mouse myeloma cells (malignant self-perpetuating antibody producing cells), and selecting out and cloning the hybrid cells (hybridomas) that produced the desired antibody reactivity. These initial monoclonal products were mouse antibodies and were very valuable in laboratory and animal research and diagnostic assays, but were problematic as therapeutic agents because of immune reactions to the foreign mouse protein. Subsequently, production of chimeric mouse-human monoclonal antibodies and means of further “humanizing” them and producing fully human recombinant monoclonal antibodies were developed. The conventions used in nomenclature of monoclonal antibodies indicate whether they are mouse derived (-omab), chimeric (-ximab), humanized (-zumab) or fully human (-umab).

Monoclonal antibodies have broad clinical and experimental medical uses. Many of the initial monoclonal antibodies used in clinical medicine were immunomodulatory agents with activity against specific immune cells, such as CD4 or CD3 lymphocytes, which are important in the pathogenesis of rejection after solid organ transplantation. Subsequently, monoclonal antibodies were prepared against specific cytokines (anti-cytokines), which were believed to play a role in cell and tissue damage in immunologically mediated diseases such as rheumatoid arthritis, ankylosing spondylitis, inflammatory bowel disease, multiple sclerosis and psoriasis, among others. In addition, therapeutic monoclonal antibodies were developed, aimed at blocking or inhibiting the activity of specific enzymes, cell surface transporters or signaling molecules and have been used in cancer chemotherapy and to treat severe viral infections. Use of monoclonal antibodies is currently broadening to therapy of other severe, nonmalignant conditions including asthma, atopic dermatitis, migraine headaches, hypercholesterolemia, osteoporosis, bacterial diseases (such as anthrax) and viral infections (such as COVID-19). Thus, the therapeutic monoclonal antibodies do not fall into a single class and have broad therapeutic uses. As of 2022, more than 80 therapeutic monoclonal antibodies have been approved for use in the United States.

Monoclonal antibodies are generally well tolerated. Because they are large proteins (typically 150-200,000 daltons in size) they require parenteral, often intravenous, administration. Circulating proteins are metabolized by many cells, but particularly by hepatocytes. Proteins undergo hepatic uptake by endocytosis and are either degraded or recycled to the cell surface for secretion. The hepatic metabolism of antibodies often determines their half-life. Proteins are broken down by cellular proteases into small peptides and amino acids that can be used to synthesize other proteins. Metabolism of proteins does not generate toxic intermediates and, therefore, monoclonal antibodies are unlikely to induce drug induced liver injury via production of toxic metabolites. On the other hand, the peptides that are generated by the metabolism of the exogenously administered protein may

ultimately be presented as foreign epitopes and generate an immune response. In addition, the primary effect of the monoclonal antibody may generate a response, either immune or otherwise, that leads to an immune mediate hepatic injury. Finally, monoclonal antibodies that suppress the immune system may cause reactivation of latent infections, including tuberculosis, herpes simplex, varicella zoster (shingles) and hepatitis B.

Among the monoclonal antibodies that have been used in clinical medicine, only a few have been linked to drug induced liver injury and, in many situations, the cause of the hepatic adverse event is often unclear. The monoclonal antibodies most clearly linked to drug induced liver injury include the antibodies to tumor necrosis factor (anti-TNF such as infliximab, adalimumab, certolizumab and golimumab) and antibodies to checkpoint proteins (anti-CTLA4 such as ipilimumab; anti-PD1 such as cemiplimab, dostarlimab, nivolumab and pembrolizumab; and anti-PD-L1 such as atezolizumab, avelumab and durvalumab) and to antibodies to B cell markers and activation signals (anti-CD20 such as rituximab, ofatumumab and tositumomab). The liver injury caused by these products is usually attributed to induction of autoimmunity or to immune modulation and reactivation of hepatitis B.

Because of their fine specificity, monoclonal antibodies can also be used to direct more conventional therapeutic agents to specific organs, tissues or cells. Several monoclonal antibody conjugates have been developed, largely in the therapy of cancer, the conjugated drug being an antineoplastic or cytotoxic agent. Five such agents are gemtuzumab ozogamicin, brentuximab vedotin, trastuzumab emtansine, inotuzumab ozogamicin and moxetumomab pasudotox. These products combine a monoclonal antibody (anti-CD33, anti-CD30, anti-HER2, anti-CD22) to a microtubule inhibitor (ozogamicin, vedotin, emtansine, pasudotox). These agents have been associated with serum enzyme elevations during therapy, and several have been linked to hepatic vascular damage, sinusoidal obstruction syndrome and nodular regenerative hyperplasia. These forms of hepatotoxicity are likely due to the conjugate (the "payload") rather than the monoclonal antibody (the targeting vehicle).

Monoclonal antibodies approved for clinical use in the United States are listed below with their brand name followed by the type of monoclonal and specific antigen targeted, the year of initial approval, likelihood score of causing liver injury, and major uses. Those monoclonal antibodies with specific chapters in LiverTox are underlined and have a link to the corresponding record. COVID-19-related monoclonal antibodies, which are typically granted Emergency Use Authorization (EUA) and are often subsequently withdrawn because of antiviral resistance, are not listed in this table but can be found in the chapter "COVID-19 Monoclonal Antibodies."

MONOCLONAL ANTIBODIES

Underlined Generic Names link to a LiverTox record.

CANCER				
Generic Name Brand Name	Type Antigenic Target	Year of Initial Approval	Likelihood Score†	Major Uses
<u>Alemtuzumab</u> Campath	Humanized CD52	2001	C	Chronic lymphocytic leukemia
<u>Atezolizumab</u> Tecentriq	Humanized PD-L1	2016	B	Urothelial carcinoma Non-small cell lung cancer Small cell lung cancer Hepatocellular carcinoma

Table continued from previous page.

Avelumab Bavencio	Human PD-L1	2017	B	Merkel cell carcinoma Urothelial carcinoma Renal cell carcinoma
Belantamab Blenrep	Humanized BCMA	2020 Withdrawn In 2022	D	Multiple myeloma
Bevacizumab Avastin	Humanized VEGF	2004	E	Colorectal cancer Renal, Ovarian, and Non-small cell lung cancer Macular degeneration (off lbl)
Blinatumomab Blincyto	Mouse CD3, CD19	2014	E*	Acute lymphoblastic leukemia
Brentuximab Adcetris	Chimeric CD30	2011	B	Hodgkin lymphoma Anaplastic large cell lymphoma
Cemiplimab Libtayo	Human PD-1	2018	C	Squamous cell carcinoma Basal cell carcinoma Non-small cell lung cancer
Cetuximab Erbix	Chimeric EGFR	2004	E	Head and neck cancer Colorectal cancer
Daratumumab Darzalex	Human CD38	2015	E	Multiple myeloma
Dinutuximab Unituxin	Chimeric GD2	2015	E*	Neuroblastoma
Dostarlimab Jemperli	Human PD-1	2021	C	Endometrial carcinoma
Durvalumab Imfinzi	Human PD-L1	2017	B	Non-small cell lung cancer Small cell lung cancer
Elotuzumab Empliciti	Humanized SLAMF7	2015	D	Multiple myeloma
Enfortumab Padcev	Nectin-4	2019	E*	Metastatic urothelial cancer
Gemtuzumab Mylotarg	Humanized CD33	2000 2017	A	Acute myelogenous leukemia
Inotuzumab Besponsa	Humanized CD22	2017	A	Acute lymphoblastic leukemia
Ipilimumab Yervoy	Human CTLA4	2009	A	Malignant melanoma Colorectal, Esophageal, Hepato-cellular and Renal cell cancer Non-small cell lung cancer
Mogamulizumab Poteligeo	Humanized CCR4	2018	C	Mycosis fungoides Sézary syndrome

Table continued from previous page.

Moxetumomab Lumoxiti	Mouse CD22	2018	E*	Hairy cell leukemia
Necitumumab Portrazza	Human EGFR	2015	E	Non-small cell lung cancer
Nivolumab Opdivo	Human PD-1	2015	A	Malignant melanoma Colorectal, Esophageal, Hepato-cellular and Renal cell cancer Non-small cell lung cancer
Ofatumumab Arzerra	Human CD20	2009	E*	Chronic lymphocytic leukemia
Olaratumab Lartruvo	Human PDGF	2016	E	Soft tissue sarcoma
Panitumumab Vectibix	Human EGFR	2006	E	Colorectal cancer
Pembrolizumab Keytruda	Humanized PD-1	2014	A	Malignant melanoma Non-small cell lung cancer Advanced cervical cancer
Pertuzumab Perjeta	Humanized HER2	2012	E	Breast cancer
Polatuzumab Vedotin Polivy	Human CD79b	2019	E*	B cell lymphomas
Ramucirumab Cyramza	Human VEGF	2014	E*	Gastric, non-small cell lung cancer Colorectal cancer
Rituximab Rituxan	Chimeric CD20	1997	A	Chronic lymphocytic leukemia Non-Hodgkin lymphoma Rheumatoid arthritis
Sacituzumab Govitecan Trodelvy	Humanized Trop-2	2020 2021 2023	E*	Triple-negative breast cancer Urothelial cancer Hormone receptor positive breast cancer
Teclistamab Tecvayli	Human BCMA, CD3	2022	D	Multiple myeloma
Tisotumab Vedotin Tivdak	Human TF	2021	E	Cervical cancer
Tositumomab Bexxar	Mouse CD20	2003 Withdrawn	E*	Non-Hodgkin lymphoma
Trastuzumab Herceptin	Humanized HER2	1998	C	Breast and gastric cancer
Trastuzumab Deruxtecan Enhertu	Humanized HER2	2019	E*	Breast, gastric, and non-small cell lung cancers
Trastuzumab Emtansine Kadcyla	Humanized HER2	2013	B	Breast cancer
Tremelimumab Imjudo	Human CTLA4	2022	C	Hepatocellular carcinoma Non-small cell lung cancer

AUTOIMMUNE DISEASES				
Generic Name Brand Name	Type Antigenic Target	Initial Approval	Likelihood Score†	Major Uses
Adalimumab Humira	Human TNF α	2002	B	Inflammatory bowel disease Rheumatoid, Psoriatic arthritis Severe psoriasis
Alemtuzumab Lemtrada	Humanized CD52	2014	C	Multiple sclerosis
Anifrolumab Saphnelo	Human Type 1 IFN	2021	E	Systemic lupus erythematosus
Belimumab Benlysta	Human B cell activity factor	2011 2020	E	Systemic lupus erythematosus Lupus nephritis
Brodalumab Siliq	Human IL-17A	2017	E	Plaque psoriasis
Canakinumab Ilaris	Human IL1 β	2009	E*	Autoinflammatory diseases
Certolizumab Cimzia	Humanized TNF α	2008	E*	Inflammatory bowel disease Rheumatoid arthritis
Daclizumab Zinbryta	Humanized CD25	2016	C	Multiple sclerosis
Dupilumab Dupixent	Human IL-4 α	2017	E	Atopic dermatitis
Efalizumab Raptiva	Humanized CD11a	2003 Withdrawn	D	Plaque psoriasis
Golimumab Simponi	Human TNF α	2009	E*	Inflammatory bowel disease Rheumatoid, Psoriatic arthritis
Guselkumab Tremfya	Human IL-23	2017	E*	Plaque psoriasis
Infliximab Remicade	Chimeric TNF α	1998	A	Inflammatory bowel disease Rheumatoid arthritis Severe psoriasis
Ixekizumab Taltz	Humanized IL-17A	2016	E	Plaque psoriasis Psoriatic arthritis
Ocrelizumab Ocrevus	Humanized CD20	2017	D	Multiple sclerosis
Omalizumab Xolair	Humanized IgE	2003	E	Eosinophilic asthma Chronic idiopathic urticaria
Risankizumab Skyrizi	Humanized IL-23	2019	E	Plaque psoriasis
Rituximab Rituxan	Chimeric CD20	1997	A	Chronic lymphocytic leukemia Non-Hodgkin lymphoma Rheumatoid arthritis
Sarilumab Kevzara	Human IL6R	2017	E*	Rheumatoid arthritis

Table continued from previous page.

Secukinumab Cosentyx	Human IL-17A	2015	E*	Plaque psoriasis Psoriatic arthritis, Ankylosing spondylitis, Scalp psoriasis Axial spondylarthritis
Siltuximab Sylvant	Chimeric IL6	2014	E	Castleman disease
Tildrakizumab Ilumya	Humanized IL-23	2018	E*	Plaque psoriasis
Tocilizumab Actemra	Humanized IL6R	2010	C	Rheumatoid arthritis Juvenile idiopathic arthritis Giant cell arteritis
Tralokinumab Adbry	Human IL-13	2021	E	Atopic dermatitis
Ustekinumab Stelara	Human IL-12, IL-23	2010	E*	Plaque psoriasis Psoriatic arthritis
Vedolizumab Entyvio	Humanized Integrin $\alpha 4\beta 7$	2014	D	Inflammatory bowel disease

LIVER TRANSPLANTATION

Generic Name Brand Name	Type Antigenic Target	Initial Approval	Likelihood Score†	Major Uses
Basiliximab Simulect	Chimeric IL-2R α	1998	E	Prevention of transplant rejection
Daclizumab Zenapax	Humanized IL-2	1997 Withdrawn	C	Prevention of transplant rejection
Muromonab-CD3 OKT3	Mouse CD3 T cells	1985 Withdrawn	E	Prevention of transplant rejection

MISCELLANEOUS

Generic Name Brand Name	Type Antigenic Target	Initial Approval	Likelihood Score†	Major Uses
Abciximab Reopro	Chimeric GpIIb/IIIa	1993		Inhibition of platelet aggregation
Aducanumab Aduhelm	Human Amyloid β	2021	E	Alzheimer disease
Alirocumab Praluent	Human PCSK9	2015	E	Hypercholesterolemia
Benralizumab Fasenra	Humanized IL5	2017	E	Eosinophilic asthma
Bezlotoxumab Zinplava	Human C. difficile toxin B	2016	E	Prevention of recurrence of C. difficile infection
Burosumab Crysvita	Human FGF 23	2018	E	X-linked Hypophosphatemia

Table continued from previous page.

Caplacizumab Cabliivi	Humanized vWF	2019	E	Acquired thrombotic thrombocytopenic purpura
Crizanlizumab Adakveo	Humanized P-selectin	2019	E	Sickle cell disease
Denosumab Prolia, Zgeva	Human RANKL	2010	E*	Osteoporosis Bone metastases
Eculizumab Soliris	Humanized C5	2007	D	Paroxysmal nocturnal hemoglobinuria
Emapalumab Gamifant	Human Interferon Gamma	2018	E	Hemophagocytic lymphohistiocytosis
Emicizumab Hemlibra	Humanized Factor IXa & X	2017	E	Hemophilia A
Eptinezumab Vyepti	Humanized CGRP	2019	E	Migraine headache
Erenumab Aimovig	Human CGRP	2018	E	Migraine headache
Evinacumab Evkeeza	Human ANGPTL3	2021	E	Hypercholesterolemia
Evolocumab Repatha	Human PCSK9	2015	E	Hypercholesterolemia
Fremanezumab Ajovy	Humanized CGRP	2018	E	Migraine headache
Galcanezumab Emgality	Humanized CGRP	2018	E	Migraine headache
Ibalizumab Trogarzo	Humanized CD4	2018	E	HIV infection
Lanadelumab Takhzyro	Human Kallikrein	2018	E	Hereditary angioneurotic edema
Lecanemab Leqembi	Human Amyloid β	2023	E	Alzheimer disease
Mepolizumab Nucala	Humanized IL15	2015	E	Eosinophilic asthma Hypereosinophilic syndrome
Natalizumab Tysabri	Humanized Integrin $\alpha 4\beta 7$	2004	B	Multiple sclerosis Inflammatory bowel disease
Obiltoximab Anthim	Chimeric Anthrax toxin	2016	E	Inhalational anthrax
Omalizumab Xolair	Humanized IgE	2003	E	Eosinophilic asthma Chronic idiopathic urticaria
Palivizumab Synagis	Humanized RSV fusion protein	1998	E	Respiratory syncytial virus infection
Ranibizumab Lucentis	Humanized VEGF-A	2006	E	Macular degeneration
Ravulizumab Ultomiris	Humanized Complement C5	2018	E*	Paroxysmal nocturnal hemoglobinuria

Table continued from previous page.

Raxibacumab	Human Anthrax toxin	2012	E	Inhalational anthrax
Reslizumab Cinqair	Humanized IL5	2016	E	Eosinophilic asthma
Romosozumab Evenity	Humanized Sclerostin	2019	E	Osteoporosis
Teplizumab Tzield	Humanized CD3	2022	E	Diabetes
Teprotumumab Tepezza	Human IGF1R	2019	E	Graves ophthalmopathy
Tezepelumab Tezspire	Human TSLP	2021	E	Asthma

† Likelihood Score indicates the likelihood of association with drug induced liver injury, based upon the known potential of the drug to cause such injury.

Updated: February 07, 2024