




Cefiderocol

Revised: October 19, 2020.

CASRN: 1225208-94-5



Drug Levels and Effects

Summary of Use during Lactation

Although no information is available on the use of cefiderocol during breastfeeding, cephalosporins are generally not expected to cause adverse effects in breastfed infants. Occasionally disruption of the infant's gastrointestinal flora, resulting in diarrhea or thrush have been reported with cephalosporins, but these effects have not been adequately evaluated. Cefiderocol is acceptable to use in nursing mothers.

Drug Levels

Maternal Levels. Relevant published information was not found as of the revision date.

Infant Levels. Relevant published information was not found as of the revision date.

Disclaimer: Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site.

Attribution Statement: LactMed is a registered trademark of the U.S. Department of Health and Human Services.

Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

Substance Identification

Substance Name

Cefiderocol

CAS Registry Number

1225208-94-5

Drug Class

Breast Feeding

Lactation

Anti-Bacterial Agents

Anti-Infective Agents

Cephalosporins