




Avapritinib

Revised: October 19, 2020.

CASRN: 1703793-34-3



Drug Levels and Effects

Summary of Use during Lactation

No information is available on the clinical use of avapritinib during breastfeeding. Because avapritinib is 98.8% bound to plasma proteins, the amounts in milk are likely to be low. However, the manufacturer recommends avoiding breastfeeding during therapy and for 2 weeks following the final dose.

Drug Levels

Maternal Levels. Relevant published information was not found as of the revision date.

Infant Levels. Relevant published information was not found as of the revision date.

Disclaimer: Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site.

Attribution Statement: LactMed is a registered trademark of the U.S. Department of Health and Human Services.

Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

Substance Identification

Substance Name

Avapritinib

CAS Registry Number

1703793-34-3

Drug Class

Breast Feeding

Lactation

Antineoplastic Agents

Enzyme Inhibitors

Protein Kinase Inhibitors

Signal Transduction Inhibitors

Tyrosine Kinase Inhibitors