



## Sucroferric Oxyhydroxide

Revised: January 15, 2024.

CASRN: 8047-67-4

### Drug Levels and Effects

#### Summary of Use during Lactation

Because sucroferric oxyhydroxide is not absorbed orally, it will not reach the bloodstream of the infant or cause any adverse effects in breastfed infants.

#### Drug Levels

*Maternal Levels.* Relevant published information was not found as of the revision date.

*Infant Levels.* Relevant published information was not found as of the revision date.

#### Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

#### Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

#### Alternate Drugs to Consider

Aluminum Hydroxide, Calcium Acetate, Calcium Carbonate, Ferric Citrate, Sevelamer

### Substance Identification

#### Substance Name

Sucroferric Oxyhydroxide

**Disclaimer:** Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site.

**Attribution Statement:** LactMed is a registered trademark of the U.S. Department of Health and Human Services.

## CAS Registry Number

8047-67-4

## Drug Class

Breast Feeding

Lactation

Milk, Human

Ferric Compounds

Iron Compounds