




Thiosulfate Sodium

Revised: September 19, 2022.

CASRN: 10102-17-7



Drug Levels and Effects

Summary of Use during Lactation

No information is available on the clinical use of sodium thiosulfate during breastfeeding. However, it is not absorbed orally and the reported side effect after oral use is diarrhea. Withholding breastfeeding for 1 to 2 hours after a dose should allow the drug to be cleared from the maternal bloodstream and breastmilk. Observe the infant for diarrhea.

Drug Levels

Maternal Levels. Relevant published information was not found as of the revision date.

Infant Levels. Relevant published information was not found as of the revision date.

Disclaimer: Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site.

Attribution Statement: LactMed is a registered trademark of the U.S. Department of Health and Human Services.

Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

Substance Identification

Substance Name

Thiosulfate Sodium

CAS Registry Number

10102-17-7

Drug Class

Breast Feeding

Lactation

Milk, Human

Antidotes

Chelating Agents