



## Teclistamab

Revised: November 15, 2023.

CASRN: 2119595-80-9

## Drug Levels and Effects

### Summary of Use during Lactation

No information is available on the clinical use of teclistamab during breastfeeding. Because teclistamab is a large protein molecule with a molecular weight of 146,000 Da, the amount in milk is likely to be very low.[1] It is also likely to be partially destroyed in the infant's gastrointestinal tract and absorption by the infant is probably minimal.[2] The manufacturer recommends that breastfeeding be withheld during therapy and for 5 months after the last dose.

### Drug Levels

*Maternal Levels.* Relevant published information was not found as of the revision date.

*Infant Levels.* Relevant published information was not found as of the revision date.

### Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

### Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

### References

1. Stratigakis A, Paty D, Zou P, et al. A regression approach for assessing large molecular drug concentration in breast milk. *Reprod Breed* 2023;3:199-207. doi:10.1016/j.repbre.2023.10.003
2. Anderson PO. Monoclonal antibodies during breastfeeding. *Breastfeed Med* 2021;16:591-3. PubMed PMID: 33956488.

**Disclaimer:** Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site.

**Attribution Statement:** LactMed is a registered trademark of the U.S. Department of Health and Human Services.

## Substance Identification

### Substance Name

Teclistamab

### CAS Registry Number

### Drug Class

Breast Feeding

Lactation

Milk, Human

Antineoplastic agents

Monoclonal antibodies