




Etrasimod

Revised: April 15, 2024.

CASRN: 1206123-37-6



Drug Levels and Effects

Summary of Use during Lactation

No information is available on the use of etrasimod during breastfeeding. Because etrasimod is 98% bound to plasma proteins, the amount in milk is likely to be low. If the mother requires etrasimod, it is not a reason to discontinue breastfeeding. Until more data become available, etrasimod should be used with caution during breastfeeding, especially while nursing a newborn or preterm infant.

Drug Levels

Maternal Levels. Relevant published information was not found as of the revision date.

Infant Levels. Relevant published information was not found as of the revision date.

Disclaimer: Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site.

Attribution Statement: LactMed is a registered trademark of the U.S. Department of Health and Human Services.

Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

Substance Identification

Substance Name

Etrasimod

CAS Registry Number

1206123-37-6

Drug Class

Breast Feeding

Lactation

Milk, Human

Gastrointestinal Agents

Sphingosine 1 Phosphate Receptor Modulators