




## Iloprost

Revised: December 15, 2023.

CASRN: 78919-13-8



## Drug Levels and Effects

### Summary of Use during Lactation

No information is available on the use of iloprost during breastfeeding. The half-life of iloprost is 20 to 30 minutes and iloprost is generally not detectable in plasma 30 to 60 minutes after inhalation. It is unlikely that clinically important amounts of iloprost are excreted into milk after 1 hour following an inhaled dose.

### Drug Levels

*Maternal Levels.* Relevant published information was not found as of the revision date.

*Infant Levels.* Relevant published information was not found as of the revision date.

**Disclaimer:** Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site.

**Attribution Statement:** LactMed is a registered trademark of the U.S. Department of Health and Human Services.

## Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

## Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

## Substance Identification

### Substance Name

Iloprost

### CAS Registry Number

78919-13-8

### Drug Class

Breast Feeding

Lactation

Milk, Human

Cardiovascular Agents

Prostaglandins, Synthetic

Vasodilator Agents