

**S13: Subgroup Analyses:** Subgroups were defined by the following attributes: (1) giving or not giving the PCP Plan before or during the wellness visit; (2) completion or not of the 2-week and 7-month surveys after the PCP visit; (3) race and ethnicity comparing white non-Hispanic individuals with DS with all others; (4) age of individuals with DS greater vs. less than or equal to 18 years, and (5) private vs. public insurance coverage. We treated loss to follow-up as wholly non-informative (i.e., completely at random) with respect to non-completion.

**Supplemental Table S4a. Subgroup analysis of primary outcome: caregivers who did versus caregivers who did not share the Primary Care Provider Plan with PCP before or during wellness visit**

Variable		Member of Caregiver Subgroup			P-value*
		Overall (N=230)	Did not share PCP Plan before wellness visit (N=13)	Did share PCP Plan before wellness visit (N=217)	
Number of indicated evaluations that were recommended or completed <sup>†</sup>	0	134 (62.0%)	7 (77.8%)	127 (61.4%)	0.267
	1	67 (31.0%)	2 (22.2%)	65 (31.4%)	
	2	13 (6.0%)	0 (0.0%)	13 (6.3%)	
	3	2 (0.9%)	0 (0.0%)	2 (1.0%)	
	[missing]	14	4	10	
Number of indicated and completed <sup>†</sup>	0	153 (70.8%)	7 (77.8%)	146 (70.5%)	0.516
	1	52 (24.1%)	2 (22.2%)	50 (24.2%)	
	2	10 (4.6%)	0 (0.0%)	10 (4.8%)	
	3	1 (0.5%)	0 (0.0%)	1 (0.5%)	
	[missing]	14	4	10	
Number of not indicated and completed	0	38 (17.6%)	1 (11.1%)	37 (17.9%)	0.184
	1	54 (25.0%)	2 (22.2%)	52 (25.1%)	
	2	56 (25.9%)	2 (22.2%)	54 (26.1%)	
	3	37 (17.1%)	1 (11.1%)	36 (17.4%)	
	4	28 (13.0%)	2 (22.2%)	26 (12.6%)	
	5	3 (1.4%)	1 (11.1%)	2 (1.0%)	
	[missing]	14	4	10	

**Notes:**

\*P-value <0.05, \*\* p-value <0.01, \*\*\* p-value <0.001

<sup>†</sup>These variables were scaled from 0–5, where each value represents the number of evaluations indicated, completed or recommended. The numbers not represented here (e.g., 4, 5) had zero responses.