

## Abstract Review Form

2. P. M. van Brussel, D. W. Eeftinck Schattenkerk, L. C. Dobrowolski, R. J. de Winter, J. A. Reekers, H. J. Verberne, L. Vogt and B. H. van den Born. Effects of renal sympathetic denervation on cardiac sympathetic activity and function in patients with therapy resistant hypertension. 2015. Int J Cardiol 202:609-614

### Attachments

PDF - 2 van Brussel, 2015.pdf ([Standard Version](#)) | ([Annotatable Version](#))

[LINK REFERENCE](#)

**BACKGROUND:** Renal sympathetic denervation (RSD) is currently being investigated in multiple studies of sympathetically driven cardiovascular diseases such as heart failure and arrhythmias. Our aim was to assess systemic and cardiac sympatholytic effects of RSD by the measurement of cardiac sympathetic activity and cardiovascular parameters.

**METHODS:** A total of 21 consecutive patients with refractory hypertension (daytime ambulatory blood pressure (BP)  $\geq 150/100$  mmHg despite the use of 3 or more antihypertensive drugs), no evidence for secondary hypertension and normal renovascular anatomy were included. RSD was performed with the Medtronic Symplicity renal denervation catheter with an average of 4.2 (range 3-6) ablations per renal artery. To assess cardiac sympathetic activity,  $^{123}\text{I}$ -mIBG cardiac scintigraphy was performed before and 6 weeks after. In addition, the effect of RSD on peripheral BP and cardiac hemodynamics were assessed non-invasively.

**RESULTS:**  $^{123}\text{I}$ -mIBG uptake before and after RSD was  $1.7 \pm 0.4\%$  vs.  $1.7 \pm 0.5\%$  at 15 min. and  $1.4 \pm 0.4\%$  vs.  $1.5 \pm 0.5\%$  after 4 h. As a consequence, washout rate was similar before ( $33.7 \pm 11.7\%$ ) and after RSD ( $30.1 \pm 12.6\%$ ,  $p=0.27$ ). In line with earlier RSD studies, a significant drop in systolic office BP ( $-12.2$  mmHg,  $p=0.04$ ) was detected, whereas the decrease in ambulatory BP was not significant. No changes were seen in heart rate, stroke volume or left ventricular contractility, both in supine position and after standing.

**CONCLUSION:** In concert with previous reports, RSD leads to a significant drop in office BP. However, a reduction in sympathetic activity could not be demonstrated on a cardiac level.

[SUBMIT FORM](#)

and go to

[This Form - Next Reference](#)

or [Skip to Next](#)

1. Exclude article because (check first that applies):

- No **original data** (e.g. review article, commentary, or editorial)
- No **human data** (e.g., evaluated outcomes in animals only)
- Not in **English**
- Does not evaluate a **renal denervation device**
- Case report** only
- Case series with **<10 patients**
- Does not apply** to any of the questions
- Other reason (specify):

[Clear Response](#)

2. Exclude, but pull for handsearching (e.g., systematic review article that applies to the question)

- Handsearch

[Clear Response](#)

3. Include or unclear so pull for full review

- Include/unclear

[Clear Response](#)

[SUBMIT FORM](#)

and go to

[This Form - Next Reference](#)

or [Skip to Next](#)