



## Luspatercept

Revised: April 15, 2023.

CASRN: 1373715-00-4

## Drug Levels and Effects

### Summary of Use during Lactation

No information is available on the clinical use of luspatercept during breastfeeding. Because luspatercept is a large protein molecule with a molecular weight of 76,000 Da, the amount in milk is likely to be very low. It is also likely to be partially destroyed in the infant's gastrointestinal tract and absorption by the infant is probably minimal. Until more data become available, the manufacturer recommends that breastfeeding be discontinued during luspatercept therapy and for 3 months after the last dose.

### Drug Levels

*Maternal Levels.* Relevant published information was not found as of the revision date.

*Infant Levels.* Relevant published information was not found as of the revision date.

### Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

### Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

## Substance Identification

### Substance Name

Luspatercept

### CAS Registry Number

1373715-00-4

**Disclaimer:** Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site.

**Attribution Statement:** LactMed is a registered trademark of the U.S. Department of Health and Human Services.

## Drug Class

Breast Feeding

Lactation

Milk, Human

Hematinics

Hematopoietic Cell Growth Factors