

Table 4. Item endorsement in percentage, by group, with the number of respondents endorsing each item in parentheses.

ITEM:	Mobility Impairment, No NB (n = 49)	NB, no UTIs (n = 47)	No mobility impairment, no NB, no UTIs (n = 64)	ALL controls (n = 160)
Cloudy Urine	4% (n = 49)	15% (n = 47)	14% (n = 64)	11% (n = 160)
Bad smelling Urine	16% (49)	15% (46)	25% (64)	19% (159)
Darker Urine	33% (49)	28% (47)	16% (64)	24% (160)
Blood in Urine	0% (49)	4% (46)	3% (64)	3% (159)
Increased Frequency of spasms	2% (49)	17% (46)	13% (64)	11% (159)
Increased urgency	41% (49)	39% (46)	49% (63)	44% (158)
Increased Catheterization	0% (1)	29% (14)	NA	27% (15)
Increased Incontinence	20% (49)	37% (46)	27% (62)	28% (157)
Increased Urine Leakage	12% (49)	27% (45)	45% (62)	29% (156)
Reduced Urine Volume during Catheterization	100% (1)	14% (14)	NA	20% (15)
Pain in Abdomen	8% (49)	11% (44)	17% (60)	12% (153)
Pain in Lower Back	20% (49)	23% (44)	52% (60)	33% (153)
Pain in Legs	41% (49)	30% (44)	32% (60)	34% (153)
Burning on Catheterization	0% (1)	21% (14)	NA	20% (15)
Burning on Passing Urine	0% (49)	2% (44)	8% (60)	4% (153)
Body Position Pain	12% (49)	23% (44)	15% (60)	16% (153)
Fever	10% (49)	7% (44)	22% (60)	14% (153)
Fatigue	39% (49)	44% (43)	38% (60)	40% (152)
Generally not Feeling Well	27% (49)	35% (43)	47% (60)	37% (152)
Muscle Aches	27% (49)	37% (43)	65% (60)	45% (152)
Altered Sleep patterns	16% (49)	28% (43)	42% (60)	30% (152)
Increased Irritability	18% (49)	23% (43)	45% (60)	30% (152)
Headaches	18% (49)	26% (43)	53% (60)	34% (152)
Dizziness	14% (49)	14% (43)	15% (60)	14% (152)
Rigor	22% (49)	35% (43)	5% (60)	19% (152)
Abdominal Bloating	8% (49)	16% (43)	31% (59)	19% (151)
Nausea	10% (49)	12% (43)	19% (59)	14% (151)
Loss of Appetite	18% (49)	9% (43)	12% (59)	13% (151)
Bowel Pattern Changes	8% (49)	16% (43)	31% (59)	19% (151)
Mean, SD total # endorsed (range)	4.6, SD = 4.3 (0–19)	5.62, SD = 5.1 (0–20)	7.0, SD = 4.4 (0–15)	5.8, SD = 4.7 (0–20)

<https://doi.org/10.1371/journal.pone.0197568.t004>