

## LETTERS

**Role of PIP<sub>2</sub> in regulating versus modulating Ca<sup>2+</sup> channel activity**

We would like to comment on an article by Gamper & Shapiro (2007) recently published in *The Journal of Physiology* entitled Target-specific PIP<sub>2</sub> signalling: How might it work? In this paper, they contend that, 'there is wide agreement that the G<sub>q/11</sub>-coupled muscarinic modulation of both N-type Ca<sup>2+</sup> and M-type K<sup>+</sup> channels is mediated by receptor-induced depletion of PIP<sub>2</sub>'. We disagree with Gamper and Shapiro's portrayal of widespread acceptance of this model for Ca<sup>2+</sup> channel modulation by G<sub>q</sub>-coupled receptors. In this paradigm, loss of PIP<sub>2</sub> association with Ca<sup>2+</sup> channels is necessary and sufficient for current inhibition (Wu *et al.* 2002; Gamper *et al.* 2004; Michailidis *et al.* 2007). Activation of phospholipase C (PLC) by G<sub>q</sub>-coupled receptors substantially lowers free PIP<sub>2</sub> levels in the plasma membrane. As a consequence, when constitutively bound PIP<sub>2</sub> dissociates and diffuses away from the channel rather than rebinding, channel activity is inhibited (Gamper *et al.* 2004). No other signalling events, e.g. kinases, phosphatases or additional lipases, participate in channel inhibition. This hypothesis arose from a similar model for M-current modulation by M<sub>1</sub>Rs, first proposed by Suh & Hille (2002) and later embraced by several labs examining Ca<sup>2+</sup> current modulation by G<sub>q</sub>-coupled receptors (Wu *et al.* 2002; Gamper *et al.* 2004; Robbins *et al.* 2006).

We have offered an alternative hypothesis to Gamper *et al.* (2004) as to the signal cascade that confers Ca<sup>2+</sup> current inhibition. Studies in our lab indicate that in addition to PLC, group IVA phospholipase A<sub>2</sub> (cPLA<sub>2</sub>) activity is required in order to observe modulation of L- and N-type Ca<sup>2+</sup> currents. Our conclusion is based on the following specific findings. (1) Arachidonic acid (AA) mimics the actions of M<sub>1</sub>R agonists in inhibiting channel activity (Liu & Rittenhouse, 2000, 2003b; Barrett *et al.* 2001; Liu *et al.* 2001, 2006). (2) The presence of bovine serum albumin (BSA), either in the pipette or in the bath solution, minimizes L- and N-current inhibition by Oxo-M (Liu & Rittenhouse, 2003b; Liu

*et al.* 2006). Moreover, AA rather than a metabolite, mediates N-current modulation since blocking AA's metabolism has no effect on exogenous AA or Oxo-M's ability to inhibit current (Barrett *et al.* 2001; Liu *et al.* 2001; Liu & Rittenhouse, 2003b). (3) Antagonizing PLA<sub>2</sub> with oleyloxyethyl phosphorylcholine (OPC) minimizes inhibition of both L- and N-currents (Liu & Rittenhouse, 2003a,b; Liu *et al.* 2004, 2006). Our OPC data contrast findings from Gamper *et al.* (2004) who reported no effect of OPC on N-current inhibition by Oxo-M. Similarly, Bannister *et al.* (2002) found that the PLA<sub>2</sub> antagonist quinacrine had no effect on M<sub>1</sub>R inhibition of recombinant L-current (Ca<sub>v</sub>1.2). However, both studies failed to provide controls demonstrating that the PLA<sub>2</sub> antagonist was effectively blocking the enzyme, so that the respective conclusions appear premature. (4) Using antibodies as functional antagonists, we found that dialysing cells with antibodies to cPLA<sub>2</sub>, but not to sPLA<sub>2</sub> or non-immunized antibody minimized L-current inhibition (Liu & Rittenhouse, 2003a; Liu *et al.* 2004, 2006). (5) Using a genetic approach we found that neurons from mice deficient in cPLA<sub>2</sub> (cPLA<sub>2</sub><sup>-/-</sup>) exhibited minimal L-current inhibition by Oxo-M (Liu *et al.* 2006). No significant differences in control current amplitude or magnitude of current inhibition by AA was observed between cPLA<sub>2</sub><sup>+/+</sup> versus cPLA<sub>2</sub><sup>-/-</sup> neurons, indicating normal channel activity in cPLA<sub>2</sub><sup>-/-</sup> neurons. Moreover, M-current inhibition by Oxo-M remained normal, indicating no change in M<sub>1</sub>R, G<sub>q</sub>, or PLC functioning in cPLA<sub>2</sub><sup>-/-</sup> neurons. However, cPLA<sub>2</sub><sup>-/-</sup> neurons exhibited decreased fatty acid release following exposure to Oxo-M compared to wild-type neurons, consistent with a requirement for cPLA<sub>2</sub>-dependent increases in free fatty acid levels in order to observe L- and N-current inhibition. (6) Using BSA as an AA scavenger to limit free fatty acid levels antagonized L- and N-current inhibition (Liu & Rittenhouse, 2003b; Liu *et al.* 2006). Taken together our findings indicate that lipid products downstream of PIP<sub>2</sub> are required for Ca<sup>2+</sup> current modulation, whereas M-current inhibition appears to occur with PIP<sub>2</sub> breakdown by PLC. Most critically, the studies with cPLA<sub>2</sub><sup>-/-</sup> neurons document that PLC activity alone is insufficient to mediate Ca<sup>2+</sup> current inhibition.

How to reconcile our previous findings that a fatty acid (probably AA) mediates Ca<sup>2+</sup> current inhibition by M<sub>1</sub>R signalling with the PIP<sub>2</sub> model remains unresolved (Liu & Rittenhouse, 2003b; Liu *et al.* 2004, 2006; Michailidis *et al.* 2007). Resolution will come with additional controls and experiments. For example the PIP<sub>2</sub> analogue DiC8-PIP<sub>2</sub> appears to minimize Ca<sup>2+</sup> current inhibition; however, this analogue of PIP<sub>2</sub> does not contain the normal fatty acid chains associated with PIP<sub>2</sub>, e.g. AA and stearic acid. It is possible that diC8-PIP<sub>2</sub> acts as a substrate competitor with other phospholipids for cPLA<sub>2</sub> decreasing the AA liberated and minimizing Ca<sup>2+</sup> channel inhibition. Similarly, whether application of exogenous phospholipids swells membranes to such an extent that M<sub>1</sub>R signalling no longer functions needs to be tested and resolved. Most critical is the need to distinguish between questions that test how PIP<sub>2</sub> functions as a regulator of channel activity *versus* its role in Ca<sup>2+</sup> channel modulation by specific G<sub>q</sub>-coupled receptors. For example, dephosphorylating PIP<sub>2</sub> may cause PIP<sub>2</sub> to dissociate from Ca<sup>2+</sup> channels and lower activity; however, during muscarinic signalling, this mechanism may play no role in decreasing current amplitude. In recent work palmitoylated charged peptides, that sequester PIP<sub>2</sub>, were dialysed into neurons, decreasing both M- and N-current amplitudes. However, low concentrations of peptide, which only minimally decreased basal current amplitude, disrupted M- but not N-current modulation by M<sub>1</sub>Rs (Robbins *et al.* 2006). The different results underscore the notion that specific experiments testing roles in modulation as well as regulation are needed in order to properly define the function of PIP<sub>2</sub> in ion channel regulation. Thus experiments outlined by Gamper & Shapiro (2007) that test how plasma membrane PIP<sub>2</sub> levels are regulated contribute to our understanding of PIP<sub>2</sub>'s functioning, but not necessarily its role in modulating Ca<sup>2+</sup> channel activity. Thus, we strongly encourage caution when using these experiments as evidence that a simple dissociation of PIP<sub>2</sub> from Ca<sup>2+</sup> channels explains how M<sub>1</sub>Rs inhibit Ca<sup>2+</sup> channel activity. Our model provides additional levels of control allowing more independent regulation of specific ion channel activity by M<sub>1</sub>R signalling.

Liwang Liu, John F. Heneghan\*  
Tora Mitra-Ganguli, Mandy L. Roberts-  
Crowley and Ann R. Rittenhouse

*Program in Neuroscience  
Department of Physiology, University  
of Massachusetts Medical School  
55 Lake Ave.  
North Worcester, MA-01655, USA*

*\*Molecular and Vascular Unit  
Department of Medicine  
Harvard medical School  
Beth Israel Deaconess Medical Center  
330 Brookline Ave.  
Boston, MA-02215, USA*

Email: liwang.lui@umassmed.edu  
jheneghan@bidmc.harvard.edu  
tora.mitra-ganguli@umassmed.edu  
mandy.roberts-crowley@umassmed.edu  
ann.rittenhouse@umassmed.edu

## References

- Bannister RA, Melliti K & Adams BA (2002). Reconstituted slow muscarinic inhibition of neuronal (Ca<sub>v</sub>1.2c), L-type Ca<sup>2+</sup> channels. *Biophys J* **32**, 56–67.
- Barrett CF, Liu L & Rittenhouse AR (2001). Arachidonic acid reversibly enhances N-type calcium current at an extracellular site. *Am J Physiol Cell Physiol* **280**, C1306–C1318.
- Gamper N, Reznikov V, Yamada Y, Yang J & Shapiro MS (2004). Phosphatidylinositol 4,5-bisphosphate signals underlie receptor-specific Gq/11-mediated modulation of N-type Ca<sup>2+</sup> channels. *J Neurosci* **24**, 10980–10992.
- Gamper N & Shapiro MS (2007). Target-specific PIP<sub>2</sub> signalling: How might it work? *J Physiol* **582**, 967–975.
- Liu L, Barrett CF & Rittenhouse AR (2001). Arachidonic acid both inhibits and enhances whole cell calcium currents in rat sympathetic neurons. *Am J Physiol Cell Physiol* **280**, C1293–C1305.
- Liu L & Rittenhouse AR (2000). Effects of arachidonic acid on unitary calcium currents in rat sympathetic neurons. *J Physiol* **525**, 391–404.
- Liu L & Rittenhouse AR (2003a). Pharmacological discrimination between muscarinic receptor signal transduction cascades with bethanechol chloride. *Br J Pharmacol* **138**, 1259–1270.
- Liu L & Rittenhouse AR (2003b). Arachidonic acid mediates muscarinic inhibition and enhancement of N-type Ca<sup>2+</sup> current in sympathetic neurons. *Proc Natl Acad Sci U S A* **100**, 295–300.
- Liu L, Roberts ML & Rittenhouse AR (2004). Phospholipid metabolism is required for M1 muscarinic inhibition of N-type calcium current in sympathetic neurons. *Eur Biophys J* **33**, 255–264.
- Liu L, Zhao R, Bai Y, Stanish LF, Evans JE, Sanderson MJ, Bonventre JV & Rittenhouse AR (2006). M1 muscarinic receptors inhibit L-type Ca<sup>2+</sup> current and M-current by divergent signal transduction cascades. *J Neurosci* **26**, 11588–11598.
- Michailidis IE, Shang Y & Yang J (2007). The lipid connection-regulation of voltage-gated Ca<sup>2+</sup> channels by phosphoinositides. *Plugs Arch* (in press).
- Robbins J, Marsh SJ & Brown DA (2006). Probing the regulation of M (Kv7) potassium channels in intact neurons with membrane-targeted peptides. *J Neurosci* **26**, 7950–7961.
- Suh BC & Hille B (2002). Recovery from muscarinic modulation of M current channels requires phosphatidylinositol 4,5-bisphosphate synthesis. *Neuron* **35**, 507–520.
- Wu L, Bauer CS, Zhen X-G, Xie C & Yang J (2002). Dual regulation of voltage-gated calcium channels by PtdIns(4,5)P<sub>2</sub>. *Nature* **419**, 947–952.