Title Page

Applying and Optimizing Water-Soluble, Slow-Release Nitrogen Fertilizers for Water-Saving Agriculture

Yanle Guo[a,b], Min Zhang[a,*], Zhiguang Liu[a,*], Chenhao Zhao[a], Hao Lu[a], Lei Zheng[c], Yuncong C. Li[d]

[a]National Engineering Laboratory for Efficient Utilization of Soil and Fertilizer Resources, National Engineering & Technology Research Center for Slow and Controlled Release Fertilizers, College of Resources and Environment, Shandong Agricultural University, Tai'an, Shandong 271018, China
[b]College of Horticulture and Landscape Architecture, Jinling Institute of Technology, Nanjing 210038, China
[c]State Key Laboratory of Nutrition Resources Integrated Utilization, Kingenta Ecological Engineering Group Co., Ltd, Linshu, Shandong 276700, China
[d]Department of Soil and Water Science, Tropical Research and Education Center, IFAS, University of Florida, Homestead, FL 33031, USA

[*] **Corresponding authors:** Tel.: +86-538-8241531; fax: +86-538-8241531
E-mail addresses: minzhang-2002@163.com (M. Zhang); liuzhiguang8235126@126.com (Z. Liu).

Table S1. The single-factor test design

Run	U:F[a] (molar ratio)	F:NH_3[b] (molar ratio)	Reaction temperature (°C)	Reaction time (h)
1	1.75	1.90	60	2.5
2	1.50	2.22	60	2.5
3	1.25	2.67	60	2.5
4	1.00	3.33	60	2.5
5	0.75	4.44	60	2.5
6	1.25	4.00	60	2.5
7	1.25	3.33	60	2.5
8	1.25	2.67	60	2.5
9	1.25	2.00	60	2.5
10	1.25	1.00	60	2.5
11	1.25	2.67	20	2.5
12	1.25	2.67	40	2.5
13	1.25	2.67	60	2.5
14	1.25	2.67	80	2.5
15	1.25	2.67	100	2.5
16	1.25	2.67	60	0.5
17	1.25	2.67	60	1.5
18	1.25	2.67	60	2.5
19	1.25	2.67	60	3.5
20	1.25	2.67	60	4.5

[a] Urea:formaldehyde ratio.

[b] Formaldehyde:amine ratio.

Figure S1. Samples of single-factor experiments

Figure S2. Samples of response surface methodology design